



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 48/00, C12N 15/63, 15/00, C07H 21/04	A1	(11) International Publication Number: WO 00/23116 (43) International Publication Date: 27 April 2000 (27.04.00)
(21) International Application Number: PCT/US99/24495 (22) International Filing Date: 19 October 1999 (19.10.99) (30) Priority Data: 60/104,994 20 October 1998 (20.10.98) US 60/125,974 24 March 1999 (24.03.99) US 09/364,862 30 July 1999 (30.07.99) US (71) Applicant (for all designated States except US): AVIGEN, INC. [US/US]; #1000, 1201 Harbor Bay Parkway, Alameda, CA 94502 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): COUTO, Linda, B. [US/US]; 7814 Oak Creek Drive, Pleasanton, CA 94588 (US). COLOSI, Peter, C. [US/US]; 2016 Encinal Avenue, Alameda, CA 94501 (US). (74) Agents: MACKNIGHT, Kamrin, T. et al.; Medlen & Carroll, LLP, Suite 2200, 220 Montgomery Street, San Francisco, CA 94104 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS (57) Abstract <p>The present invention provides improved viral vectors useful for the expression of genes at high levels in human cells. In particular, the present invention provides adeno-associated vectors (AAV) suitable for gene therapy. These vectors are capable of delivering nucleic acid containing constructs which result in the production of full-length therapeutic levels of biologically active Factor VIII in the recipient individual <i>in vivo</i>. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using these constructs.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII BY TARGET CELLS

This application claims priority benefit of U.S. provisional application Nos.
5 60/125,974 filed March 24, 1999 and 60/104,994 filed October 20, 1998, which are
hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to AAV vectors suitable for hemophilia gene
10 therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids
encoding Factor VIII into a recipient subject suffering from hemophilia A, such that
the subject's blood is able to clot.

BACKGROUND

15 Hemophilia is a genetic disease characterized by a blood clotting deficiency. In
hemophilia A (classic hemophilia, Factor VIII deficiency), an X-chromosome-linked
genetic defect disrupts the gene encoding Factor VIII, a plasma glycoprotein, which is
a key component in the blood clotting cascade. Human Factor VIII is synthesized as a
single chain polypeptide, with a predicted molecular weight of 265 kDa. The Factor
20 VIII gene codes for 2351 amino acids, and the protein has six domains, designated
from the amino to the carboxy terminus as A1-A2-B-A3-C1-C2 (Wood *et al.*, Nature
312:330 [1984]; Vehar *et al.*, Nature 312:337 [1984]; and Toole *et al.*, Nature 312:342
[1984]). Human Factor VIII is processed within the cell to yield a heterodimer
primarily comprised of a heavy chain of 200 kDa containing the A1, A2, and B
25 domains and an 80 kDa light chain containing the A3, C1, and C2 domains (Kaufman
et al., J. Biol. Chem., 263:6352-6362 [1988]). Both the single chain polypeptide and
the heterodimer circulate in the plasma as inactive precursors (Ganz *et al.*, Eur. J.
Biochem., 170:521-528 [1988]). Activation of Factor VIII in plasma is initiated by
thrombin cleavage between the A2 and B domains, which releases the B domain and
30 results in a heavy chain consisting of the A1 and A2 domains. The 980 amino acid B

domain is deleted in the activated procoagulant form of the protein. Additionally, in the native protein, two polypeptide chains ("a" and "b"), flanking the B domain, are bound to a divalent calcium cation. Hemophilia may result from point mutations, deletions, or mutations resulting in a stop codon (*See, Antonarakis et al., Mol. Biol. Med., 4:81 [1987]*).

The disease is relatively rare, afflicting approximately one in 10,000 males. Hemophilia in females is extremely rare, although it may occur in female children of an affected father and carrier mother, as well as in females with X-chromosomal abnormalities (*e.g., Turner syndrome, X mosaicism, etc.*). The severity of each patient's disease is broadly characterized into three groups--"mild," "moderate," and "severe," depending on the severity of the patient's symptoms and circulating Factor VIII levels. While normal levels of Factor VIII range between 50 and 200 ng/mL plasma, mildly affected patients have 6-60% of this value, and moderately affected patients have 1-5% of this value. Severely affected hemophiliacs have less than 1% of normal Factor VIII levels.

While hemophiliacs clearly require clotting factor after surgery or severe trauma, on a daily basis, spontaneous internal bleeding is a greater concern. Hemophiliacs experience spontaneous hemorrhages from early infancy, as well as frequent spontaneous hemarthroses and other hemorrhages requiring clotting factor replacement. Without effective treatment, chronic hemophilic arthropathy occurs by young adulthood. Severely affected patients are prone to serious hemorrhages that may dissect through tissue planes, ultimately resulting in death due to compromised vital organs.

Hematomas are commonly observed in moderately and severely affected hemophiliacs. In these patients, hematomas have a tendency to progressively enlarge and dissect in all directions. Some of these hematomas expand locally, resulting in local compression of adjacent organs, blood vessels, and nerves. A rare, yet often fatal, complication of abdominal hematomas is the perforation and drainage of the hematoma into the colon, resulting in infection and septicemia. Intracranial and/or extracranial hemorrhage also represent very dangerous bleeding situations. While

subcutaneous hematomas may dissect into muscle, pharyngeal and retropharyngeal hematomas (e.g., complicating bacterial or viral pharyngitis) may enlarge and obstruct the airway, sometimes resulting in a life-threatening situation that requires administration of a sufficient dose of Factor VIII concentrate to normalize the Factor VIII level.

In addition to hematomas, hemarthroses are commonly observed in hemophiliacs, with bleeding into the joint accounting for approximately 75% of hemophilic bleeding. Repeated hemorrhaging into the joints eventually results in extensive destruction of articular cartilage, synovial hyperplasia, and other reactive changes in adjacent tissues and bone. A major complication of repeated hemarthroses is joint deformity, which is often accompanied by muscle atrophy and soft tissue contractures; osteoporosis and cystic areas in the subchondral bone may also develop, along with progressive loss of joint space.

Other symptoms are often observed in hemophiliacs, including hematuria and mucous membrane bleeding. Hematuria is experienced by virtually all severely affected hemophiliacs sometime during their lifetimes, and mucous membrane bleeding is common in hemophiliacs. Bone cysts (pseudotumors) are rare, but dangerous complications of hemophilic bleeding. In many of these cases, immediate treatment is necessary.

In the early 1980s, many severely affected hemophiliacs were treated with Factor VIII concentrate about three times weekly. Unfortunately, these concentrates transmitted viruses, such as hepatitis B and/or C, and human immunodeficiency virus (HIV). In the United States and Western Europe, at least 75% of Factor VIII concentrate recipients have been reported to have anti-HIV antibodies (*See*, Schrier and Leung, *supra*). Some of these patients also developed HIV-associated immune thrombocytopenia, a very serious complication in hemophiliacs. In spite of antiviral therapy (e.g., with zidovudine and pentamidine prophylaxis), which has tended to slow disease progression, full-blown AIDS (acquired immunodeficiency syndrome) occurs at an inexorable rate in hemophiliacs infected with HIV. Indeed, this has reversed the improvement in the life expectancy of hemophiliacs, which peaked at 66 years of age

during the 1970s, and has dropped to 49 years (*See, Schrier and Leung, supra*). The development of virus-free preparations and recombinant Factor VIII has helped control infectious viral contamination.

However, for hemophiliacs, the availability of viral-free concentrates and recombinant Factor VIII, while significant, is but part of the solution. In order to prevent spontaneous internal bleeding episodes, patients suffering from hemophilia A must consistently have serum Factor VIII levels of about 1%, and preferably 5%. Currently, the cost of viral-free concentrates and recombinant Factor VIII make it prohibitively expensive to administer the clotting factor prophylactically or on a maintenance basis. Indeed, most hemophiliacs in the U.S. do not receive recombinant Factor VIII therapy on a maintenance basis, but only receive it prior to activities or events which might cause bleeding (*e.g.*, surgery), or as a treatment for spontaneous bleeding.

Moreover, even if cost effective preparations of recombinant or virus-free Factor VIII were available, a steady state level of Factor VIII cannot be achieved by its daily administration. At best, patients receive widely varying levels of Factor VIII. Immediately following the administration, the levels are super-physiological, while prior to administration the levels are sub-physiological. Thus, there remains a need for methods and compositions that are relatively economic, yet effective in the treatment and prevention of bleeding in hemophiliacs, particularly spontaneous bleeds. Furthermore, there is a need in the art for methods and compositions for long term delivery of clotting factors (*e.g.*, Factor VIII) which more closely mimic the steady state physiological levels observed in normal individuals.

SUMMARY OF THE INVENTION

The present invention provides improved viral vectors suitable for gene therapy to treat hemophilia. In particular, the present invention provides AAV vectors and methods for treating hemophilia A by delivering nucleic acids coding for the clotting protein Factor VIII. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the vectors.

The present invention is particularly suited for use in hemophilia A gene therapy. Accordingly, in one embodiment of the invention, at least one AAV vector containing a nucleic acid molecule encoding Factor VIII is operably linked to control sequences that direct expression of Factor VIII in a suitable recipient cell. The AAV
5 vectors are then introduced into a recipient cell of the subject, under conditions that result in expression of Factor VIII. The subject, therefore, has a continuous supply of Factor VIII available to clot blood during bleeding episodes.

Using the methods of the present invention, long term expression of therapeutic levels of Factor VIII have been achieved *in vivo*. In one embodiment, animals were
10 administered, via the portal vein, two AAV vectors: one carrying the DNA sequence coding for the heavy chain of Factor VIII and the other carrying the DNA sequence coding for the light chain of Factor VIII. Blood samples were collected periodically and assayed for Factor VIII activity. Reproducibly, animals expressed between 600 and 900 ng/ml of biologically active Factor VIII, levels that are well above the normal
15 physiological levels of approximately 200 ng/ml. Furthermore, these levels have been sustained for over 13 months without a decrease in Factor VIII levels or activity. In a related embodiment, a B-domain deleted form of Factor VIII was cloned into a single AAV vector and shown to express biologically active Factor VIII.

It is not intended, however, that the present invention be limited to specific
20 embodiments. Many different forms of recombinant Factor VIII have been made and tested both *in vitro* and *in vivo*, using a variety of different control and regulatory sequences. Any DNA sequence coding for biologically active Factor VIII can be expressed using the AAV vectors and methods taught in the present invention. Therefore, the present invention encompasses any AAV vector or vectors containing
25 Factor VIII sequences that produce biologically active Factor VIII protein *in vitro* or *in vivo*.

For example, in some embodiments, the AAV vector contains the first 57 base pairs of the Factor VIII heavy chain which encodes the 10 amino acid signal sequence, as well as the human growth hormone (hGH) polyadenylation sequence. In some
30 alternative embodiments, the vector also contains the A1 and A2 domains, as well as 5

amino acids from the N-terminus of the B domain, and/or 85 amino acids of the C-terminus of the B domain, as well as the A3, C1, and C2 domains. In yet other embodiments, the nucleic acids coding for Factor VIII heavy chain and light chain were cloned into a single vector separated by 42 nucleic acids coding for 14 amino acids of the B domain.

The present invention also provides methods for administering the above-described vectors. For example, it is intended that the present invention encompass methods suitable for delivery of the AAV vectors to the livers of recipient patients or test animals. It is not intended that the present invention be limited to any particular route of administration. However, in preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal or arterial vasculature.

These and other embodiments of the invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a schematic representation of the Factor VIII protein.

Figure 2 provides a schematic representation of a B-domain deleted form of Factor VIII protein.

Figure 3 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (AAV-F8-1) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 4 provides a schematic representation of a B-domain deleted Factor VIII AAV construct (PVM4.1c-F8AB) from internal terminal repeat (ITR to ITR), including control sequences.

Figure 5 provides the sequence of pAAV-F8-1 (ITR to ITR), with the plasmid backbone omitted.

Figure 6 provides the sequence of pVm4.1cF8ΔB (ITR to ITR), with the plasmid backbone omitted.

Figure 7 provides a map of rAAV-hFVIII-HC and rAAV-hFVIII-LC vectors.

Figure 8 provides a graph demonstrating the expression of various human FVIII constructs in mouse plasma.

Figure 9 provides Southern blot analyses of liver DNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

5 Figure 10 provides Southern blot analyses of DNA from different tissues using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

Figure 11 provides Northern blot analyses of liver RNA using probes specific for (A) the light chain of hFVIII, and (B) the heavy chain of hFVIII.

10 GENERAL DESCRIPTION OF THE INVENTION

The present invention relates to improved viral vectors useful for expressing gene products at high levels in human cells. In particular, the present invention provides AAV vectors suitable for gene therapy. These vectors are capable of delivering nucleic acid containing constructs which result in the production of Factor
15 VIII protein in a host. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

The AAV vectors and rAAV virions of the present invention can be produced using standard methodology known to those of skill in the art. The methods generally
20 involve the steps of: (1) introducing an AAV vector into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell, wherein the helper virus and/or
25 accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions. The AAV vector, AAV helper construct and the helper virus or accessory function vector(s) can be introduced into the host cell either simultaneously or serially, using standard transfection techniques.

Unless otherwise indicated, the practice of the present invention employs conventional methods of virology, microbiology, molecular biology and recombinant DNA techniques within the skill of the art, including those described in such references as Sambrook *et al.* (eds.) *Molecular Cloning: A Laboratory Manual*; Glover
5 (ed.) *DNA Cloning: A Practical Approach*, Vols. I and II; Gait (ed.) *Oligonucleotide Synthesis*; Hames and Higgins (eds.) *Nucleic Acid Hybridization*; Hames and Higgins (eds.) *Transcription and Translation*; Tijessen (ed.) *CRC Handbook of Parvoviruses*, Vols. I and II; and Fields and Knipe (eds.) *Fundamental Virology*, 2nd Edition, Vols. I and II.

10

Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

As used herein, the terms "gene transfer" and "gene delivery" refer to methods
15 or systems for reliably inserting a particular nucleotide sequence (*e.g.*, DNA) into targeted cells. In particularly preferred embodiments, the nucleotide sequence comprises at least a portion of Factor VIII.

As used herein, the terms "vector," and "gene transfer vector" refer to any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus,
20 virion, etc., which is capable of replication when associated with the proper control sequences and/or which can transfer nucleic acid sequences between cells. Thus, the term includes cloning and expression vectors, as well as viral vectors.

Gene transfer vectors may include transcription sequences such as polyadenylation sites, selectable markers or reporter genes, enhancer sequences, and
25 other control sequences which allow for the induction of transcription. Such control sequences are described more fully below.

The term "expression vector" as used herein refers to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular

host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, as well as other sequences. Eukaryotic cells are generally known to utilize promoters (constitutive, inducible or tissue specific), enhancers, and termination and polyadenylation signals, although some elements may be deleted and other elements added without sacrificing the necessary expression.

As used herein, the terms "host" and "expression host" refer to organisms and/or cells which harbor an exogenous DNA sequence (e.g., via transfection), an expression vector or vehicle, as well as organisms and/or cells that are suitable for use in expressing a recombinant gene or protein. It is not intended that the present invention be limited to any particular type of cell or organism. Indeed, it is contemplated that any suitable organism and/or cell will find use in the present invention as a host.

As used herein, the terms "viral replicons" and "viral origins of replication" refer to viral DNA sequences that allow for the extrachromosomal replication of a vector in a host cell expressing the appropriate replication factors. In some embodiments, vectors which contain either the SV40 or polyoma virus origin of replication replicate to high copy number, while vectors which contain the replicons from bovine papillomavirus or Epstein-Barr virus replicate extrachromosomally at low copy number may be utilized in other embodiments.

As used herein, the term "AAV vector" refers to a vector having functional or partly functional ITR sequences. The ITR sequences may be derived from an adeno-associated virus serotype, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. The ITRs, however, need not be the wild-type nucleotide sequences, and may be altered (e.g., by the insertion, deletion or substitution of nucleotides), so long as the sequences retain function provide for functional rescue, replication and packaging. AAV vectors can have one or more of the AAV wild-type genes deleted in whole or part, preferably the *rep* and/or *cap* genes but retain functional flanking ITR sequences. Functional ITR sequences are necessary for the rescue, replication and packaging of the AAV virion. Thus, an "AAV vector" is

defined herein to include at least those sequences required in *cis* for replication and packaging (e.g., functional ITRs) of the virus.

As used herein, the term "ITR" refers to inverted terminal repeats. The terms "adeno-associated virus inverted terminal repeats" or "AAV ITRs" refer to the art-recognized palindromic regions found at each end of the AAV genome which function together in *cis* as origins of DNA replication and as packaging signals for the virus. For use in some embodiments of the present invention, flanking AAV ITRs are positioned 5' and 3' of one or more selected heterologous nucleotide sequences. Optionally, the ITRs together with the *rep* coding region or the Rep expression product provide for the integration of the selected sequences into the genome of a target cell.

As used herein, the term "AAV *rep* coding region" refers to the art-recognized region of the AAV genome which encodes the replication proteins Rep 78, Rep 68, Rep 52 and Rep 40. These Rep expression products have been shown to possess many functions, including recognition, binding and nicking of the AAV origin of DNA replication, DNA helicase activity and modulation of transcription from AAV (or other heterologous) promoters. The Rep expression products are collectively required for replicating the AAV genome. Muzyczka (Muzyczka, Curr. Top. Microbiol. Immunol., 158:97-129 [1992]) and Kotin (Kotin, Hum. Gene Ther., 5:793-801 [1994]) provide additional descriptions of the AAV *rep* coding region, as well as the *cap* coding region described below. Suitable homologues of the AAV *rep* coding region include the human herpesvirus 6 (HHV-6) *rep* gene which is also known to mediate AAV-2 DNA replication (Thomson *et al.*, Virol., 204:304-311 [1994]).

As used herein, the term "AAV *cap* coding region" refers to the art-recognized region of the AAV genome which encodes the capsid proteins VP1, VP2, and VP3, or functional homologues thereof. These *cap* expression products supply the packaging functions which are collectively required for packaging the viral genome.

As used herein, the term "AAV helper function" refers to AAV coding regions capable of being expressed in the host cell to complement AAV viral functions missing from the AAV vector. Typically, the AAV helper functions include the AAV

rep coding region and the AAV *cap* coding region. An "AAV helper construct" is a vector containing AAV coding regions required to complement AAV viral functions missing from the AAV vector (*e.g.*, the AAV *rep* coding region and the AAV *cap* coding region).

5 As used herein, the terms "accessory functions" and "accessory factors" refer to functions and factors that are required by AAV for replication, but are not provided by the AAV virion (or rAAV virion) itself. Thus, these accessory functions and factors must be provided by the host cell, a virus (*e.g.*, adenovirus or herpes simplex virus), or another expression vector that is co-expressed in the same cell. Generally, the E1,
10 E2A, E4 and VA coding regions of adenovirus are used to supply the necessary accessory function required for AAV replication and packaging (Matsushita *et al.*, Gene Therapy 5:938 [1998]).

As used herein, the term "wild type" ("wt") refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a
15 naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product which displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that
20 naturally-occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

As used herein, the term "AAV virion" refers to a complete virus particle, such as a "wild-type" (wt) AAV virus particle (comprising a linear, single-stranded AAV nucleic acid genome associated with an AAV capsid protein coat). In this regard,
25 single-stranded AAV nucleic acid molecules of either complementary sense (*e.g.*, "sense" or "antisense" strands), can be packaged into any one AAV virion and both strands are equally infectious.

As used herein, the terms "recombinant AAV virion," and "rAAV virion" refer to an infectious viral particle containing a heterologous DNA molecule of interest (*e.g.*,

Factor VIII sequence) which is flanked on both sides by AAV ITRs. In some embodiments of the present invention, an rAAV virion is produced in a suitable host cell which contains an AAV vector, AAV helper functions and accessory functions introduced therein. In this manner, the host cell is rendered capable of encoding AAV polypeptides that are required for packaging the AAV vector containing a recombinant nucleotide sequence of interest, such as at least a portion of Factor VIII or portions of Factor VIII domains, into recombinant virion particles for subsequent gene delivery.

As used herein, the term "transfection" refers to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art (*See e.g.*, Graham *et al.*, *Virol.*, 52:456 [1973]; Sambrook *et al.*, *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratories, New York [1989]; Davis *et al.*, *Basic Methods in Molecular Biology*, Elsevier, [1986]; and Chu *et al.*, *Gene* 13:197 [1981]. Such techniques can be used to introduce one or more exogenous DNA moieties, such as a gene transfer vector and other nucleic acid molecules, into suitable recipient cells.

As used herein, the terms "stable transfection" and "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated foreign DNA into the genomic DNA.

As used herein, the term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells which have taken up foreign DNA but have failed to integrate this DNA.

As used herein, the term "transduction" denotes the delivery of a DNA molecule to a recipient cell either *in vivo* or *in vitro*, via a replication-defective viral vector, such as via a recombinant AAV virion.

As used herein, the term "recipient cell" refers to a cell which has been transfected or transduced, or is capable of being transfected or transduced, by a nucleic acid construct or vector bearing a selected nucleotide sequence of interest (*i.e.*, Factor VIII). The term includes the progeny of the parent cell, whether or not the progeny are identical in morphology or in genetic make-up to the original parent, so long as the selected nucleotide sequence is present.

The term "heterologous" as it relates to nucleic acid sequences such as coding sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence in nature. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (*e.g.*, synthetic sequences having codons different from the native gene). Similarly, a cell transfected with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention. Allelic variation or naturally occurring mutational events do not give rise to heterologous DNA, as used herein.

As used herein, "coding sequence" or a sequence which "encodes" a particular antigen, is a nucleic acid sequence which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo*, when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the coding sequence.

As used herein, the term "nucleic acid" sequence refers to a DNA or RNA sequence. The term captures sequences that include any of the known base analogues

of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl)uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 5 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 10 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

15 As used herein, the term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

As used herein, the term "regulatory element" refers to a genetic element which controls some aspect of the expression of nucleic acid sequences. For example, a 20 promoter is a regulatory element which facilitates the initiation of transcription of an operably linked coding region. Other regulatory elements are splicing signals, polyadenylation signals, termination signals, etc. (defined *infra*).

The term DNA "control sequences" refers collectively to regulatory elements such as promoter sequences, polyadenylation signals, transcription termination 25 sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control sequences need always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate recipient cell.

Transcriptional control signals in eukaryotes generally comprise "promoter" and "enhancer" elements. Promoters and enhancers consist of short arrays of DNA sequences that interact specifically with cellular proteins involved in transcription (Maniatis *et al.*, Science 236:1237 [1987]). Promoter and enhancer elements have been isolated from a variety of eukaryotic sources including genes in yeast, insect and mammalian cells and viruses (analogous control sequences, *i.e.*, promoters, are also found in prokaryotes). The selection of a particular promoter and enhancer depends on what cell type is to be used to express the protein of interest (*i.e.*, Factor VIII). Some eukaryotic promoters and enhancers have a broad host range while others are functional in a limited subset of cell types (*See e.g.*, Voss *et al.*, Trends Biochem. Sci., 11:287 [1986]; and Maniatis *et al.*, *supra*, for reviews). For example, the SV40 early gene enhancer is very active in a wide variety of cell types from many mammalian species and has been widely used for the expression of proteins in mammalian cells (Dijkema *et al.*, EMBO J. 4:761 [1985]). Two other examples of promoter and enhancer elements active in a broad range of mammalian cell types are those from the human elongation factor 1 α gene (Uetsuki *et al.*, J. Biol. Chem., 264:5791 [1989]; Kim *et al.*, *Gene* 91:217 [1990]; and Mizushima and Nagata, Nucl. Acids. Res., 18:5322 [1990]) and the long terminal repeats of the Rous sarcoma virus (Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79:6777 [1982]) and the human cytomegalovirus (Boshart *et al.*, Cell 41:521 [1985]). Promoters and enhancers can be found naturally alone or together. For example, the long terminal repeats of retroviruses contain both promoter and enhancer functions. Moreover, generally promoters and enhancers act independently of the gene being transcribed or translated. Thus, the enhancer and promoter may be "endogenous" or "exogenous" or "heterologous." An "endogenous" enhancer/promoter is one which is naturally linked with a given gene in the genome. An "exogenous" or "heterologous" enhancer and promoter is one which is placed in juxtaposition to a gene by means of genetic manipulation (*i.e.*, molecular biological techniques) such that transcription of that gene is directed by the linked enhancer/promoter.

As used herein, the term "tissue specific" refers to regulatory elements or control sequences, such as a promoter, enhancers, etc., wherein the expression of the nucleic acid sequence is substantially greater in a specific cell type(s) or tissue(s). In particularly preferred embodiments, the albumin promoter and the transthyretin promoter display increased expression of FVIII in hepatocytes, as compared to other cell types. It is not intended, however, that the present invention be limited to the albumin or transthyretin promoters or to hepatic-specific expression, as other tissue specific regulatory elements, or regulatory elements that display altered gene expression patterns, are contemplated.

The presence of "splicing signals" on an expression vector often results in higher levels of expression of the recombinant transcript. Splicing signals mediate the removal of introns from the primary RNA transcript and consist of a splice donor and acceptor site (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York [1989], pp. 16.7-16.8). A commonly used splice donor and acceptor site is the splice junction from the 16S RNA of SV40.

Efficient expression of recombinant DNA sequences in eukaryotic cells requires expression of signals directing the efficient termination and polyadenylation of the resulting transcript. Transcription termination signals are generally found downstream of the polyadenylation signal and are a few hundred nucleotides in length. The term "poly A site" or "poly A sequence" as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript. Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. The poly A signal utilized in an expression vector may be "heterologous" or "endogenous." An endogenous poly A signal is one that is found naturally at the 3' end of the coding region of a given gene in the genome. A heterologous poly A signal is one which is one which is isolated from one gene and placed 3' of another gene. A commonly used heterologous poly A signal is the SV40 poly A signal. The SV40 poly A signal is contained on a 237 bp *Bam*HI/*Bcl*II restriction fragment and directs both termination and polyadenylation (Sambrook *et al.*, *supra*, at 16.6-16.7).

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control sequences operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control sequences need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

The term "isolated" when used in relation to a nucleic acid, as in "an isolated oligonucleotide" or "isolated polynucleotide" refers to a nucleic acid sequence that is identified and separated from at least one contaminant nucleic acid with which it is ordinarily associated in its natural source. Isolated nucleic acid is such present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids are nucleic acids such as DNA and RNA found in the state they exist in nature. For example, a given DNA sequence (*e.g.*, a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs which encode a multitude of proteins. The isolated nucleic acid, oligonucleotide, or polynucleotide may be present in single-stranded or double-stranded form. When an isolated nucleic acid, oligonucleotide or polynucleotide is to be utilized to express a protein, the oligonucleotide or polynucleotide will contain at a minimum the sense or coding strand (*i.e.*, the oligonucleotide or polynucleotide may single-stranded), but may contain both the sense and anti-sense strands (*i.e.*, the oligonucleotide or polynucleotide may be double-stranded).

As used herein, the term "purified" or "to purify" refers to the removal of contaminants from a sample. For example, antibodies may be purified by removal of contaminating non-immunoglobulin proteins; they may also purified by the removal of immunoglobulin that does not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII). The removal of non-immunoglobulin proteins and/or the removal of

immunoglobulins that do not bind the antigen of interest (*e.g.*, at least a portion of Factor VIII) results in an increase in the percent of desired antigen-reactive immunoglobulins in the sample. In another example, recombinant polypeptides of Factor VIII are expressed in bacterial host cells and the polypeptides are purified by the removal of host cell proteins; the percent of recombinant polypeptides is thereby
5 increased in the sample.

As used herein, the term "chimeric protein" refers to two or more coding sequences obtained from different genes, that have been cloned together and that, after translation, act as a single polypeptide sequence. Chimeric proteins are also referred to as "hybrid proteins." As used herein, the term "chimeric protein" refers to coding
10 sequences that are obtained from different species of organisms, as well as coding sequences that are obtained from the same species of organisms.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequence. The
15 composition may comprise an aqueous solution.

As used herein, the term "at risk" is used in references to individuals who are at risk for experiencing hemorrhagic episodes. In particularly preferred embodiments, the individuals are hemophiliacs with mild, moderate, or severe hemophilia.

As used herein, the term "subject" refers to any animal (*i.e.*, vertebrates and
20 invertebrates), while the term "vertebrate subject" refers to any member of the subphylum Chordata. It is intended that the term encompass any member of this subphylum, including, but not limited to humans and other primates, rodents (*e.g.*, mice, rats, and guinea pigs), lagamorphs (*e.g.*, rabbits), bovines (*e.g.*, cattle), ovines (*e.g.*, sheep), caprines (*e.g.*, goats), porcines (*e.g.*, swine), equines (*e.g.*, horses),
25 canines (*e.g.*, dogs), felines (*e.g.*, cats), domestic fowl (*e.g.*, chickens, turkeys, ducks, geese, other gallinaceous birds, etc.), as well as feral or wild animals, including, but not limited to, such animals as ungulates (*e.g.*, deer), bear, fish, lagamorphs, rodents, birds, etc. It is not intended that the term be limited to a particular age or sex. Thus,

adult and newborn subjects, as well as fetuses, whether male or female, are encompassed by the term.

As defined herein, a "therapeutically effective amount" or "therapeutic effective dose" is an amount or dose of AAV vector or virions capable of producing sufficient amounts of Factor VIII to decrease the time it takes for a subject's blood to clot. Generally, severe hemophiliacs having less than 1% of normal levels of FVIII have a whole blood clotting time of greater than 60 minutes as compared to approximately 10 minutes for non-hemophiliacs.

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to AAV vectors suitable for hemophilia A gene therapy. More particularly, these AAV vectors are suitable for delivering nucleic acids encoding Factor VIII into a recipient host suspected of suffering from a blood clotting disorder. Using the nucleic acid as a template, the host produces Factor VIII, such that the subject's blood is able to clot. The present invention also provides pharmaceutical compositions comprising such AAV vectors, as well as methods for making and using the constructs.

I. AAV Vectors

Adeno-associated virus (AAV) is a non-pathogenic, replication-defective, helper-dependent parvovirus (or "dependovirus" or "adeno-satellite virus"). There are at least six recognized serotypes, designated as AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-X7, etc. Culture and serologic evidence indicates that human infection occurs with AAV-2 and AAV-3. Although 85% of the human population is seropositive for AAV-2, the virus has never been associated with disease in humans. Recombinant AAV (rAAV) virions are of interest as vectors for gene therapy because of their broad host range, excellent safety profile, and duration of transgene expression in infected hosts. One remarkable feature of recombinant AAV (rAAV) virions is the prolonged expression achieved after *in vivo* administration.

AAV vectors of the present invention may be constructed using known techniques to provide, as operatively linked components in the direction of transcription, (a) control sequences including a transcriptional initiation and termination regions, and (b) a nucleotide sequence encoding at least a portion of Factor VIII. The control sequences are selected to be functional in a targeted recipient cell. The resulting construct which contains the operatively linked components is bounded (5' and 3') with functional AAV ITR sequences.

The nucleotide sequences of AAV ITR regions are known (*See e.g.*, Kotin, Hum. Gene Ther., 5:793-801 [1994]; Berns, "Parvoviridae and Their Replication" in Fields and Knipe (eds), *Fundamental Virology*, 2nd Edition, for the AAV-2 sequence). AAV ITRs used in the vectors of the invention need not have a wild-type nucleotide sequence, and may be altered (*e.g.*, by the insertion, deletion or substitution of nucleotides). Additionally, AAV ITRs may be derived from any of several AAV serotypes, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX7, etc. Furthermore, 5' and 3' ITRs which flank a selected nucleotide sequence in an AAV vector need not necessarily be identical or derived from the same AAV serotype or isolate, so long as they function as intended.

A. Control Sequences

In some embodiments of the present invention, heterologous control sequences are employed with the vectors. Useful heterologous control sequences generally include those derived from sequences encoding mammalian or viral genes. Examples include, but are not limited to, the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter (Ad MLP), a herpes simplex virus (HSV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter region (CMVIE), a rous sarcoma virus (RSV) promoter, synthetic promoters, hybrid promoters, and the like. In addition, sequences derived from nonviral genes, such as the murine metallothionein gene, also find use herein. Such promoter sequences are commercially available (*e.g.*, from Stratagene).

It is contemplated that in some embodiments, tissue-specific expression may be desirable (e.g., expression of biologically active Factor VIII by hepatocytes). It is not intended that the present invention be limited to expression of biologically active Factor VIII by any particular cells or cell types. However, as hepatocytes (*i.e.*, liver cells) are the cells that normally synthesized Factor VIII (*See*, Kaufman, *Ann. Rev. Med.*, 43:325 [1992]), it is contemplated that in some particularly preferred embodiments, the compositions of the present invention be administered to the liver.

In preferred embodiments, expression is achieved by coupling the coding sequence for Factor VIII with heterologous control sequences derived from genes that are specifically transcribed by a selected tissue type. A number of tissue-specific promoters have been described above which enable directed expression in selected tissue types. However, control sequences used in the present AAV vectors can also comprise control sequences normally associated with the selected nucleic acid sequences.

B. Construction of AAV Factor VIII Vectors

AAV vectors that contain a control sequence and a nucleotide sequence of interest (*i.e.*, at least a portion of the sequence encoding Factor VIII), bounded by AAV ITRs (*i.e.*, AAV vectors), can be constructed by directly inserting selected sequences into an AAV genome with the major AAV open reading frames ("ORFs") excised. Other portions of the AAV genome can also be deleted, so long as a sufficient portion of the ITRs remain to allow for replication and packaging functions. These constructs can be designed using techniques well known in the art (*See e.g.*, U.S. Patent Nos. 5,173,414 and 5,139,941, all of which are herein incorporated by reference); International Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski *et al.*, *Mol. Cell. Biol.*, 8:3988-3996 [1988]; Vincent *et al.*, *Vaccines 90* [Cold Spring Harbor Laboratory Press, 1990]; Carter, *Curr. Opin. Biotechnol.*, 3:533-539 [1992]; Muzyczka, *Curr. Top. Microbiol. Immunol.*, 158:97-129 [1992]; Kotin, *Hum. Gene*

Ther., 5:793-801 [1994]; Shelling and Smith, *Gene Ther.*, 1:165-169 [1994]; and Zhou *et al.*, *J. Exp. Med.*, 179:1867-1875 [1994]).

Alternatively, AAV ITRs can be excised from the viral genome or from an AAV vector containing the same and fused 5' and 3' of a selected nucleic acid construct that is present in another vector using standard ligation techniques, such as those described in Sambrook *et al.*, *supra*. For example, ligations can be accomplished in 20 mM Tris-Cl pH 7.5, 10 mM MgCl₂, 10 mM DTT, 33 µg/ml BSA, 10 mM-50 mM NaCl, and either 40 µM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0°C (for "sticky end" ligation) or 1 mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14°C (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 30-100 µg/ml total DNA concentrations (5-100 nM total end concentration). AAV vectors which contain ITRs have been described in (*e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference). In particular, several AAV vectors are described therein which are available from the American Type Culture Collection ("ATCC") under Accession Numbers 53222, 53223, 53224, 53225 and 53226.

Additionally, chimeric genes can be produced synthetically to include AAV ITR sequences arranged 5' and 3' of a selected nucleic acid sequence. The complete chimeric sequence is assembled from overlapping oligonucleotides prepared by standard methods (*See e.g.*, Edge, *Nature* 292:756 [1981]; Nambair *et al.*, *Science* 223:1299 [1984]; and Jay *et al.*, *J. Biol. Chem.*, 259:6311 [1984]).

Moreover, it is not intended that the present invention be limited to any specific Factor VIII sequence. Many natural and recombinant forms of Factor VIII have been isolated and assayed both *in vitro* and *in vivo*, using a variety of different regulatory elements and control sequences. Therefore, any known, or later discovered, DNA sequence coding for biologically active Factor VIII can be expressed, alone or in combination with at least one additional vector, using the AAV vectors and methods taught in the present invention. Examples of naturally occurring and recombinant forms of Factor VIII can be found in the patent and scientific literature including, U.S. 5,563,045, U.S. 5,451,521, U.S. 5,422,260, U.S. 5,004,803, U.S. 4,757,006, U.S.

5,661,008, U.S. 5,789,203, U.S. 5,681,746, U.S. 5,595,886, U.S. 5,045,455, U.S. 5,668,108, U.S. 5,633,150, U.S. 5,693,499, U.S. 5,587,310, U.S. 5,171,844, U.S. 5,149,637, U.S. 5,112,950, U.S. 4,886,876, WO 94/11503, WO 87/07144, WO 92/16557, WO 91/09122, WO 97/03195, WO 96/21035, WO 91/07490, EP 0 672 138, EP 0 270 618, EP 0 182 448, EP 0 162 067, EP 0 786 474, EP 0 533 862, EP 0 506 757, EP 0 874 057, EP 0 795 021, EP 0 670 332, EP 0 500 734, EP 0 232 112, EP 0 160 457, Sanberg *et al.*, XXth Int. Congress of the World Fed. Of Hemophilia (1992), and Lind *et al.*, Eur. J. Biochem., 232:19 (1995).

Nucleic acid sequences coding for the above-described Factor VIII can be obtained using recombinant methods, such as by screening cDNA and genomic libraries from cells expressing Factor VIII or by deriving the sequence from a vector known to include the same. Furthermore, the desired sequence can be isolated directly from cells and tissues containing the same, using standard techniques, such as phenol extraction and PCR of cDNA or genomic DNA (*See e.g.*, Sambrook *et al.*, *supra*, for a description of techniques used to obtain and isolate DNA). Nucleotide sequences encoding an antigen of interest (*i.e.*, Factor VIII sequence) can also be produced synthetically, rather than cloned. The complete sequence can be assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence (*See e.g.*, Edge, Nature 292:756 [1981]; Nambair *et al.*, Science 223:1299 [1984]; and Jay *et al.*, J. Biol. Chem., 259:6311 [1984]).

Although it is not intended that the present invention be limited to any particular methods for assessing the production of biologically active Factor VIII, such methods as immunoassays (*e.g.*, ELISA) and biological activity assays are contemplated (*e.g.*, coagulation activity assays).

Furthermore, while in particularly preferred embodiments, human Factor VIII is encompassed by the present invention, it is not intended that the present invention be limited to human Factor VIII. Indeed, it is intended that the present invention encompass Factor VIII from animals other than humans, including but not limited to companion animals (*e.g.*, canines, felines, and equines), livestock (*e.g.*, bovines,

caprines, and ovines), laboratory animals (e.g., rodents such as murines, as well as lagamorphs), and "exotic" animals (e.g., marine mammals, large cats, etc.).

II. Virion Production

5 Producing AAV Factor VIII vectors and rAAV Factor VIII virions of the present invention generally involve the steps of: (1) introducing an AAV vector containing the Factor VIII gene into a host cell; (2) introducing an AAV helper construct into the host cell, where the helper construct includes AAV coding regions capable of being expressed in the host cell to complement AAV helper functions
10 missing from the AAV vector; (3) introducing one or more helper viruses and/or accessory function vectors into the host cell, wherein the helper virus and/or accessory function vectors provide accessory functions capable of supporting efficient recombinant AAV ("rAAV") virion production in the host cell; and (4) culturing the host cell to produce rAAV virions.

15 The above-described vectors and constructs can be introduced into a cell using standard methodology known to those of skill in the art (e.g., transfection). A number of transfection techniques are generally known in the art (*See e.g.*, Graham *et al.*, Virol., 52:456 [1973], Sambrook *et al. supra*, Davis *et al., supra*, and Chu *et al.*, Gene 13:197 [1981]). Particularly suitable transfection methods include calcium phosphate
20 co-precipitation (Graham *et al.*, Virol., 52:456-467 [1973]), direct micro-injection into cultured cells (Capecchi, Cell 22:479-488 [1980]), electroporation (Shigekawa *et al.*, BioTechn., 6:742-751 [1988]), liposome-mediated gene transfer (Mannino *et al.*, BioTechn., 6:682-690 [1988]), lipid-mediated transduction (Felgner *et al.*, Proc. Natl. Acad. Sci. USA 84:7413-7417 [1987]), and nucleic acid delivery using high-velocity
25 microprojectiles (Klein *et al.*, Nature 327:70-73 [1987]).

 For the purposes of the invention, suitable host cells for producing rAAV virions include microorganisms, yeast cells, insect cells, and mammalian cells, that can be, or have been, used as recipients of a heterologous DNA molecule. The term includes the progeny of the original cell which has been transfected. Thus, as

indicated above, a "host cell" as used herein generally refers to a cell which has been transfected with an exogenous DNA sequence. Cells from the stable human cell line, 293 (ATCC Accession No. CRL1573) are preferred in the practice of the present invention. Particularly, the human cell line 293 is a human embryonic kidney cell line
5 that has been transformed with adenovirus type-5 DNA fragments (Graham *et al.*, J. Gen. Virol., 36:59 [1977]), and expresses the adenoviral E1a and E1b genes (Aiello *et al.*, Virol., 94:460 [1979]). The 293 cell line is readily transfected, and provides a particularly convenient platform in which to produce rAAV virions.

Host cells containing the above-described AAV vectors must be rendered
10 capable of providing AAV helper functions in order to replicate and encapsidate the nucleotide sequences flanked by the AAV ITRs to produce rAAV virions. AAV helper functions are generally AAV-derived coding sequences which can be expressed to provide AAV gene products that, in turn, function in *trans* for productive AAV replication. AAV helper functions are used herein to complement necessary AAV
15 functions that are missing from the AAV vectors. Thus, AAV helper functions include one, or both of the major AAV ORFs, namely the *rep* and *cap* coding regions, or functional homologues thereof.

AAV helper functions are introduced into the host cell by transfecting the host cell with an AAV helper construct either prior to, or concurrently with, the transfection
20 of the AAV vector. AAV helper constructs are thus used to provide at least transient expression of AAV *rep* and/or *cap* genes to complement missing AAV functions that are necessary for productive AAV infection. AAV helper constructs lack AAV ITRs and can neither replicate nor package themselves.

In preferred embodiments, these constructs are in the form of a vector,
25 including, but not limited to, plasmids, phages, transposons, cosmids, viruses, or virions. A number of AAV helper constructs have been described, such as the commonly used plasmids pAAV/Ad and pIM29+45 which encode both Rep and Cap expression products (See *e.g.*, Samulski *et al.*, J. Virol., 63:3822-3828 [1989]; and McCarty *et al.*, J. Virol., 65:2936-2945 [1991]). A number of other vectors have been

described which encode Rep and/or Cap expression products (*See e.g.*, U.S. Patent No. 5,139,941, herein incorporated by reference).

Both AAV vectors and AAV helper constructs can be constructed to contain one or more optional selectable markers. Suitable markers include genes which confer
5 antibiotic resistance or sensitivity to, impart color to, or change the antigenic characteristics of those cells which have been transfected with a nucleic acid construct containing the selectable marker when the cells are grown in an appropriate selective medium. Several selectable marker genes that are useful in the practice of the invention include the gene encoding aminoglycoside phosphotransferase (APH) that
10 allows selection in mammalian cells by conferring resistance to G418 (Sigma). Other suitable markers are known to those of skill in the art.

The host cell (or packaging cell) must also be rendered capable of providing non-AAV derived functions, or "accessory functions," in order to produce rAAV virions. Accessory functions are non-AAV derived viral and/or cellular functions upon
15 which AAV is dependent for its replication. Thus, accessory functions include at least those non-AAV proteins and RNAs that are required in AAV replication, including those involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of *rep* and *cap* expression products and AAV capsid assembly. Viral-based accessory functions can be derived from any of
20 the known helper viruses.

Particularly, accessory functions can be introduced into and then expressed in host cells using methods known to those of skill in the art. Commonly, accessory functions are provided by infection of the host cells with an unrelated helper virus. A number of suitable helper viruses are known, including adenoviruses; herpesviruses
25 such as herpes simplex virus types 1 and 2; and vaccinia viruses. Nonviral accessory functions will also find use herein, such as those provided by cell synchronization using any of various known agents (*See e.g.*, Buller *et al.*, J. Virol., 40:241-247 [1981]; McPherson *et al.*, Virol., 147:217-222 [1985]; and Schlehofer *et al.*, Virol., 152:110-117 [1986]).

Alternatively, accessory functions can be provided using an accessory function vector. Accessory function vectors include nucleotide sequences that provide one or more accessory functions. An accessory function vector is capable of being introduced into a suitable host cell in order to support efficient AAV virion production in the host cell. Accessory function vectors can be in the form of a plasmid, phage, virus, transposon or cosmid. Accessory vectors can also be in the form of one or more linearized DNA or RNA fragments which, when associated with the appropriate control sequences and enzymes, can be transcribed or expressed in a host cell to provide accessory functions.

Nucleic acid sequences providing the accessory functions can be obtained from natural sources, such as from the genome of adenovirus, or constructed using recombinant or synthetic methods known in the art. In this regard, adenovirus-derived accessory functions have been widely studied, and a number of adenovirus genes involved in accessory functions have been identified and partially characterized (*See e.g., Carter, "Adeno-Associated Virus Helper Functions," in CRC Handbook of Parvoviruses*, Vol. I (P. Tijssen, ed.) [1990], and Muzyczka, *Curr. Top. Microbiol. Immun.*, 158:97-129 [1992]). Specifically, early adenoviral gene regions E1a, E2a, E4, VAI RNA and, possibly, E1b are thought to participate in the accessory process (Janik *et al.*, *Proc. Natl. Acad. Sci. USA* 78:1925-1929 [1981]). Herpesvirus-derived accessory functions have been described (*See e.g., Young et al., Prog. Med. Virol.*, 25:113 [1979]). Vaccinia virus-derived accessory functions have also been described (*See e.g., Carter, supra., and Schlehofer et al., Virol.*, 152:110-117 [1986]).

As a consequence of the infection of the host cell with a helper virus, or transfection of the host cell with an accessory function vector, accessory functions are expressed which transactivate the AAV helper construct to produce AAV Rep and/or Cap proteins. The Rep expression products direct excision of the recombinant DNA (including the DNA of interest encoding at least a portion of Factor VIII) from the AAV vector. The Rep proteins also serve to duplicate the AAV genome. The expressed Cap proteins assemble into capsids, and the recombinant AAV genome is

packaged into the capsids. Thus, productive AAV replication ensues, and the DNA is packaged into rAAV virions.

Following recombinant AAV replication, rAAV virions can be purified from the host cell using a variety of conventional purification methods, such as CsCl
5 gradients. Further, if helper virus infection is employed to express the accessory functions, residual helper virus can be inactivated, using known methods. For example, adenovirus can be inactivated by heating to temperatures of approximately 60°C for approximately 20 minutes or more, as appropriate. This treatment selectively inactivates the helper adenovirus which is heat labile, while preserving the rAAV
10 which is heat stable.

III. Pharmaceutical Compositions

The resulting rAAV virions are then ready for use in pharmaceutical compositions which can be delivered to a subject, so as to allow production of
15 biologically active Factor VIII. Pharmaceutical compositions comprise sufficient genetic material that allows the recipient to produce a therapeutically effective amount of Factor VIII so as to reduce, stop and/or prevent hemorrhage. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible
20 pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, clotting factors or factor precursors, drugs or hormones. In preferred embodiments, the pharmaceutical compositions also contain a pharmaceutically acceptable excipient. Such excipients include any pharmaceutical
25 agent that does not itself induce an immune response harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, glycerol, sugars and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides,
30 hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such

as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. A thorough discussion of pharmaceutically acceptable excipients is available in *Remington's Pharmaceutical Sciences* (Mack Pub. Co., N.J. [1991]).

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran.

Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art (e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes).

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment. For administration of Factor VIII-containing vectors, such labeling would include amount, frequency, and method of administration.

5 Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. Determining a therapeutic effective dose is well within the capability of those skilled in the art using the techniques taught in the present invention, such as ELISA and ChromZ FVIII coagulation activity assay, and other
10 techniques known in the art. Therapeutic doses will depend on, among other factors, the age and general condition of the subject, the severity of hemophilia, and the strength of the control sequences. Thus, a therapeutically effective amount in humans will fall in a relatively broad range that can be determined through clinical trials.

It is intended that the dosage treatment and regimen used with the present
15 invention will vary, depending upon the subject and the preparation to be used. Thus, the dosage treatment may be a single dose schedule or a multiple dose schedule. Moreover, the subject may be administered as many doses as appropriate to achieve or maintain the desired blood clotting time.

Direct delivery of the pharmaceutical compositions *in vivo* will generally be
20 accomplished via injection using a conventional syringe, although other delivery methods such as convention-enhanced delivery are envisioned (*See e.g.*, United States Patent No. 5,720,720, incorporated herein by reference). In this regard, the compositions can be delivered subcutaneously, epidermally, intradermally, intrathecally, intraorbitally, intramucosally (*e.g.*, nasally, rectally and vaginally),
25 intraperitoneally, intravenously, intraarterially, orally, or intramuscularly. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal applications. In particularly preferred embodiments, the compositions are administered intravenously in the portal vasculature or hepatic artery

One skilled in the art will recognize that the methods and compositions
30 described above are also applicable to a range of other treatment regimens known in

the art. For example, the methods and compositions of the present invention are compatible with *ex vivo* therapy (e.g., where cells are removed from the body, incubated with the AAV vector and the treated cells are returned to the body).

5 IV. ADMINISTRATION

AAV vector can be administered to any tissue suitable for the expression of Factor VIII. In a preferred embodiments, the AAV vectors of the present invention are successfully administered via the portal vasculature or hepatic artery where it is thought, without being bound by theory, that the vector transduces hepatocytes.

10 Current approaches to targeting genes to the liver have focused upon *ex vivo* gene therapy. *Ex vivo* liver-directed gene therapy involves the surgical removal of liver cells, transduction of the liver cells *in vitro* (e.g., infection of the explanted cells with recombinant retroviral vectors) followed by injection of the genetically modified liver cells into the liver or spleen of the patient. A serious drawback for *ex vivo* gene
15 therapy of the liver is the fact that hepatocytes cannot be maintained and expanded in culture. Therefore, the success of *ex vivo* liver-directed gene therapy depends upon the ability to efficiently and stably engraft the genetically modified (*i.e.*, transduced) hepatocytes and their progeny. It has been reported that even under optimal conditions, autologous modified liver cells injected into the liver or spleen which
20 engraft represent only a small percentage (less than 10%) of the total number of cells in the liver. Ectopic engraftment of transduced primary hepatocytes into the peritoneal cavity has been tried, in order to address the problem of engraftment in the liver.

Given the problems associated with *ex vivo* liver-directed gene therapy, *in vivo* approaches have been investigated for the transfer of genes into hepatocytes, including
25 the use of recombinant retroviruses, recombinant adenoviruses, liposomes and molecular conjugates. While these *in vivo* approaches do not suffer from the drawbacks associated with *ex vivo* liver-directed gene therapy, they do not provide a means to specifically target hepatocytes. In addition, several of these approaches require performance of a partial hepatectomy, in order to achieve prolonged expression

of the transferred genes *in vivo*. Adenovirus and molecular conjugate based delivery methods also result in liver toxicity and inflammation which is an undesirable feature of Factor VIII gene therapy. The present invention provides compositions and methods for the long-term expression of biologically active Factor VIII. It is contemplated that the present invention will bypass the need for partial hepatectomy, while allowing expression of Factor VIII in concentrations that are therapeutic *in vivo*. The present invention further provides gene therapy compositions and methods that target hepatocytes for the production of Factor VIII by treated individuals.

Other tissues, however, may be suitable for the expression of Factor VIII even if they are not the tissue that normally synthesizes the protein. Muscle cells, for example, have been shown to express biologically active blood clotting Factor IX even though it is normally synthesized in the liver.

Finally, the AAV vectors may contain any nucleic acid sequences coding for biologically active Factor VIII. Additionally, the AAV vectors may contain a nucleic acid coding for fragments of Factor VIII which is itself not biologically active, yet when administered into the subject improves or restores the blood clotting time. For example, as discussed above, the Factor VIII protein comprises two polypeptide chains: a heavy chain and a light chain separated by a B-domain which is cleaved during processing. As demonstrated by the present invention, co-transducing recipient cells with the Factor VIII heavy and light chains leads to the expression of biologically active Factor VIII. Because, however, most hemophiliacs contain a mutation or deletion in only one of the chains (*e.g.*, heavy or light chain), it may be possible to administer only the chain defective in the patient and allow the patient to supply the other chain. In this case, the AAV vector would fall within the scope of the invention even though the single chain (*i.e.*, heavy or light) would not be biologically active until it was administered into a subject which can supply the second chain, thus forming biologically active Factor VIII.

V. FACTOR VIII ASSAYS

As described in the Experimental section below, there are many ways to assay Factor VIII expression and activity. Although the present invention is not limited to immunoassay methods, the present invention also provides methods for detecting Factor VIII expression comprising the steps of: a) providing a sample suspected of containing Factor VIII, and a control containing a known amount of known Factor VIII; and b) comparing the test sample with the known control, to determine the relative concentration of Factor VIII in the sample. Thus, the methods are capable of identifying samples (*e.g.*, patient samples) with sufficient or insufficient quantities of Factor VIII. In addition, the methods may be conducted using any suitable means to determine the relative concentration of Factor VIII in the test and control samples, including but not limited to means selected from the group consisting of Western blot analysis, Northern blot analysis, Southern blot analysis, denaturing polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), reverse transcriptase-coupled polymerase chain reaction (RT-PCR), enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescent immunoassay (IFA). Thus, the methods may be conducted to determine the presence of normal Factor VIII sequences in the genome of the animal source of the test sample, or the expression of Factor VIII (mRNA or protein), as well as detect the presence of abnormal or mutated Factor VIII gene sequences in the test samples.

In one preferred embodiment, the presence of Factor VIII is detected by immunochemical analysis. For example, the immunochemical analysis can comprise detecting binding of an antibody specific for an epitope of Factor VIII. In one another preferred embodiment of the method, the antibody comprises polyclonal antibodies, while in another preferred embodiment, the antibody comprises monoclonal antibodies.

It is further contemplated that antibodies directed against at least a portion of Factor VIII will be used in methods known in the art relating to the localization and structure of Factor VIII (*e.g.*, for Western blotting), measuring levels thereof in appropriate biological samples, etc. The antibodies can be used to detect Factor VIII in a biological sample from an individual (*e.g.*, an individual treated using the methods

and/or compositions of the present invention). The biological sample can be a biological fluid, including, but not limited to, blood, serum, plasma, interstitial fluid, urine, cerebrospinal fluid, synovial fluid, and the like. In particular, the antigen can be detected from cellular sources, including, but not limited to, hepatocytes. For
5 example, cells can be obtained from an individual and lysed (*e.g.*, by freeze-thaw cycling, or treatment with a mild cytolytic detergent including, but not limited to, TRITON X-100, digitonin, NONIDET P (NP)-40, saponin, and the like, or combinations thereof; *See, e.g.*, International Patent Publication WO 92/08981).

The biological samples can then be tested directly for the presence of the
10 Factor VIII using an appropriate strategy (*e.g.*, ELISA or RIA) and format (*e.g.*, microwells, dipstick [*e.g.*, as described in International Patent Publication WO 93/03367], etc.). Alternatively, proteins in the sample can be size separated (*e.g.*, by polyacrylamide gel electrophoresis (PAGE), with or without sodium dodecyl sulfate (SDS), and the presence of Factor VIII detected by immunoblotting [*e.g.*, Western
15 blotting]). Immunoblotting techniques are generally more effective with antibodies generated against a peptide corresponding to an epitope of a protein, and hence, are particularly suited to the present invention. In another preferred embodiment, the level of Factor VIII is assayed using the whole-blood clotting time and activated partial thromboplastin time (aPTT) of the subject's blood using techniques well known in the
20 art (Herzog *et al.*, Nature Medicine 5:56 [1999]).

The foregoing explanations of particular assay systems are presented herein for purposes of illustration only, in fulfillment of the duty to present an enabling disclosure of the invention. It is to be understood that the present invention contemplates a variety of immunochemical assay protocols within its spirit and scope.
25 Indeed, other methods such as biological assays to determine the presence and activity of Factor VIII are also encompassed by the present invention.

Thus, in addition to the immunoassay systems described above, other assay systems, such as those designed to measure and/or detect Fraction VIII and/or clotting

ability of a subject's blood are also encompassed by the present invention (e.g., the ChromZ FVIII coagulation activity [FVIII-c] assay [Helena Labs]).

EXPERIMENTAL

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

In the experimental disclosure which follows, the following abbreviations apply: N (Normal); M (Molar); mM (millimolar); μ M (micromolar); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); l or L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nm (nanometers); mU (milliunits); ^{51}Cr (Chromium 51); μCi (microcurie); EC (degrees Centigrade); hFVIII (human factor VIII); FVIII (factor VIII); pH (hydrogen ion concentration); JRH grade; NaCl (sodium chloride); HCl (hydrochloric acid); OD (optical density); bp (base pair(s)); ATP (adenosine 5'-triphosphate); PCR (polymerase chain reaction); DNA (deoxyribonucleic acid); cDNA (complementary DNA); AAV (adeno-associated virus); rAAV (recombinant adeno-associated virus); ITR (inverted terminal repeat); FCS or FBS (fetal calf serum; fetal bovine serum); CFA (complete Freund's adjuvant); BSA (bovine serum albumin); ATCC (American Type Culture Collection, Rockville, MD); Sigma (Sigma Aldrich, St. Louis, MO); Biodesign International (Biodesign International, Kennebunk, MI); Baxter Hyland (Baxter Healthcare Corp., Biotech Group--Hyland Division, Hayward, CA); Helena Labs (Helena Laboratories, Beaumont, TX); American Diagnostica (American Diagnostica, Greenwich, CT); Accurate Chemical (Accurate Chemical and Scientific Corp., Westbury, NY); Molecular Probes (Molecular Probes, Eugene, OR); Vysis (Vysis, Downer Grove, IL); Tel-Test (Tel-Test, Inc., Friendswood, TX); Molecular Dynamics (Molecular Dynamics, Sunnyvale, CA); NUNC (Naperville, IL); and Stratagene

(Stratagene Cloning Systems, La Jolla, CA); and Biodesign (Biodesign International, Kennebunkport, ME).

EXAMPLE 1

Dual Vector Plasmid Construction

5 The heavy and light chains of human Factor VIII (hFVIII) were assembled according to those reported by Yonemura *et al* (Yonemura *et al.*, Prot. Engineer., 6:669-674 [1993]) and cloned as expression cassettes into AAV vectors. Both vectors contain the promoter and the first non-coding intron (from -573 to +985) from the
10 human elongation factor 1 α (EF1 α) gene (Uetsuki *et al*, J. Biol. Chem., 264:5791-5798 [1989]; and Kim *et al.*, Gene 9:217-223 [1990]). Each vector also contains the first 57 base pairs of the FVIII heavy chain encoding the 19 amino acid signal sequence. The heavy chain construct encodes the A1 and A2 domains and 5 amino acids from the N terminus of the B domain. The light chain vector encodes 85 amino
15 acids of the carboxy terminal B domain, in addition to the A3, C1, and C2 domains. Both vectors utilize the human growth hormone (hGH) polyadenylation signal. The expression cassettes were inserted between AAV ITRs. The initial cloning step involved deleting 854 bp of EF1 α sequences between the *SpeI* and *XcmI* sites of pVm4.1e-hFIX (Nakai *et al.*, Blood 91:1-9 [1998]), and religating to create
20 pVm4.1e δ D-hFIX.

This construct was then digested with *EcoRI*, which released the hFIX cDNA, and was ligated to an oligonucleotide containing *MfeI* ends (*EcoRI*-compatible) and an internal *ClaI* restriction site, creating pVm4.1e δ D-linker. The heavy and light chain fragments, including the hGH polyadenylation sequences were isolated from
25 pVm4.1cFVIII-HC and pVm4.1cFVIII-LC, respectively as *ClaI*-*BstEII* fragments. These fragments were cloned between the corresponding sites in the pVm4.1e δ D-linker, creating plasmids pVm4.1e δ D-FVIII-HC (also, rAAV-hFVIII-HC) and pVm4.1e δ D-FVIII-LC (also, rAAV-hFVIII-LC).

Figure 7 provides a map of the constructs. In this figure, the upper line in each panel represents the gene structure of the vectors, and the lower line represents the structure of the hFVIII protein domains encoded by the vectors (ITR, AAV inverted terminal repeat; EF1 α Pro/Intron 1, human polypeptide elongation factor 1 α gene promoter and first intron; hFVIII-HC human FVIII cDNA; hFVIII-LC, human FVIII cDNA; hGH PA, human growth hormone polyadenylation signal; SS, human FVIII signal sequence; A1, A2, "B", A3, C1, C2, complete and incomplete ("") protein domains of the hFVIII protein).

EXAMPLE 2

Single Vector Plasmid Construction

The plasmid pAAV-F8-1 construct containing both the light and heavy chains of factor VIII was constructed as follows. A PCR fragment, Z8, containing cloning sites, 5' -splicing donor site of a synthetic intron based on EF1 α and immunoglobulin G (IgG) intron sequences, Kozak sequence and the first 16 nucleotides of the human blood coagulation factor VIII (FVIII) coding sequence was generated using oligonucleotides Z8S and Z8A. The sequences of the nucleic acids is shown below:

Oligonucleotide Z8S:

20 5' cccaagcttgccggccgcccgggtgccgcccctaggcaggttaagtccgtgtgtggttcc 3'
(SEQ ID NO:1)

Oligonucleotide Z8A:

25 5' ccgctcgagcagagctctatttgcattggtggaatcgatgccgcccgaaccacacacggc 3'
(SEQ ID NO:2)

PCR fragment Z8:

30 5' cccaagcttgccggccgcccgggtgccgcccctaggcaggttaagtccgtgtgtggttccgc
ggcatcgattccaccatgcaaataagagctctgctcgagcgg 3' (SEQ ID NO:3)

Nucleic acid Z8 was inserted into pZERO-2 (Invitrogen) between *HindIII* and *XhoI* sites to create pZ8. A PCR fragment, INT3, containing the branching point, the

polypyrimidine tract, and the 3' splicing acceptor site of the synthetic intron was generated using oligonucleotides INT3S and INT3A whose sequence is shown below.

Oligonucleotide INT3S:

5 5' ttccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactga 3'
 (SEQ ID NO:4)

Oligonucleotide INT3A:

 5' gaatcgataacctgtggagaaaaagaaaaagtggatgtcagtgtagtaattcaaggc 3'
 10 (SEQ ID NO:5)

PCR Fragment INT3:

 5' ttccgcgggcctggcctctttacgggttatggcccttgcgtgccttgaattactgacact
 gacatccactttttctttctccacaggtatcgattc 3' (SEQ ID NO:6)
 15

 INT3 was inserted between the *Sac*II and *Cla*I sites of pZ8 to create pZ8.I. Therefore, Z8.I contains the entire synthetic intron between *Avr*II and *Cla*I sites. A hFVIII cDNA fragment having *Sac*I and *Xho*I restriction sites was inserted between the *Sac*I and *Xho*I sites of pZ8.I to create pZ8.I.dB. Therefore, pZ8.I.dB contains a
 20 synthetic intron and the entire coding sequence of hFVIII.

 A PCR fragment, EG3, containing three HNF-3 binding sites and -54 to +8 of mouse albumin gene was generated using oligonucleotides EG3S and EG3A with modifications to eliminate linker sequences. The sequences of EG3S and EG3A are as follows:

25

Oligonucleotide EG3S:

 5' aggggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaataccatccaggggaatgtttgttcttaaatacca
 tctacagttattggttaaa 3' (SEQ ID NO:7)

30

Oligonucleotide EG3A:

5' ggaaagtgatctgtgtgcagaaagactcgctctaataacttctttaaccaataactg 3'

(SEQ ID NO:8)

5 PCR Fragment EG3:

5' agggaaatgtttgttcttaaataccatccagggaatgtttgttcttaaataccatccagggaatgtttgttctta
aataccatctacagttattggtaaagaagtattagagcgagctcttctgcacacagatcacctttcc 3'

(SEQ ID NO:9)

10 EG3 was then phosphorylated using T4 polynucleotide kinase and inserted into the *Sma*I site of pZ8.I.dB to create pZ8.I.dB.egg. A DNA fragment, SPA, containing an efficient synthetic polyA signal based on rabbit β -globin sequences (Genes and Develop., 3:1019) was generated by hybridizing two oligonucleotides SPA.S and SPA.A.

15

Oligonucleotide SPA.S:

5' tcgagaataaaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgc 3'

(SEQ ID NO:10)

20 Oligonucleotide SPA.A:

5' tcgagcgccgcacacaaaaaccaacacacagatctctagagctctgatcttttattc 3'

(SEQ ID NO:11)

PCR Fragment SPA:

25 5' tcgagaataaaaagatcagagctctagagatctgtgtgttggtttttgtgtgcggccgctcga 3'

(SEQ ID NO:12)

SPA was inserted into the *Xho*I site of pZero-2 to create pZero-2.SPA. SPA was excised from a pZero-2.SPA clone and inserted into the *Xho*I site of pZ8.I.dB.egg

to create pZ8.I.dB.egg.A. pAAV-CMV-FIX9 was digested with *Cla*I, blunted with T4 polymerase and religated to create pAAV(*Cla*-)-CMV-FIX9.

The entire expression cassette containing HNF-3.albumin promoter-synthetic intron-hFVIII-synthetic poly A signal was excised from pZ8.I.dB.egg.A using *Not*I and
5 ligated to the plasmid backbone and AAV ITRs from pAAV (*Cla*-)-CMV-FIX9 to create pAAV-F8-1. The nucleotide sequence of the vector from ITR to ITR (*i.e.*, excluding plasmid backbone) is shown in SEQ ID NO 13.

EXAMPLE 3

Virion Production

10

AAV vectors were produced from these plasmids using the Ad free system as previously described in U.S. Patent No. 5,858,351; U.S. Patent No. 5,846,528; U.S. Patent No. 5,622,856; and Matsushita *et al.*, Gene Ther 5:938 (1998) all of which are hereby incorporated by reference. Briefly, 293 cells (ATCC, catalog number CRL-
15 1573) were seeded in 10 cm dishes at a density of 3×10^6 cells per dish in 10 ml medium and incubated at 37°C with CO₂ and humidity. After an overnight incubation, cells were approximately seventy to eighty percent confluent.

The cells were then transfected with DNA by the calcium phosphate method, as is well known in the art. Briefly, 7 µg of AAV vector containing the Factor VIII
20 coding region, 7 µg of pladenos which supplies the accessory functions, and 7 µg of 1909 AAV helper were added to a 3 ml sterile, polystyrene snap cap tube using sterile pipette tips. Then, 1.0 ml of 300 mM CaCl₂ (JRH grade) was added to each tube and mixed by pipetting up and down. An equal volume of 2X HBS (274 mM NaCl, 10 mM KCl, 42 mM HEPES, 1.4 mM Na₂PO₄, 12 mM dextrose, pH 7.05, JRH grade)
25 was added with a 2 ml pipette, and the solution was pipetted up and down three times. The DNA mixture was immediately added to the cells, one drop at a time, evenly throughout the 10 cm dish. The cells were then incubated at 37°C with CO₂ and humidity for six hours. A granular precipitate was visible in the transfected cell cultures. After six hours, the DNA mixture was removed from the cells, which were
30 provided with fresh medium and incubated for 72 hours.

After 72 hours, the cells were harvested, pelleted, and resuspended in 1 ml TBS/1% BSA. Freeze/thaw extracts were prepared by repeatedly (three times) freezing the cell suspension on dry ice and thawing at 37°C. Viral preps were stored at -80°C and titered by dot blot assay prior to the first round of infection.

5

EXAMPLE 4

In Vitro Cell Transduction

Cells from the stable human cell line, 293 (ATCC No. CRL1573) were seeded in six-well plates (*i.e.*, plates having six wells for cell growth) at a density of 5×10^5 cells/well. When the monolayers reached 80-90% confluence, they were infected with rAAV virions AAV-e δ D-FVIII-HC, AAV-e δ D-FVIII-LC, an equal ratio of AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC, or AAV-e δ D-FIX at MOIs of 3×10^3 and 3×10^4 . Eighteen hours post infection, the media were replaced with DMEM/10% heat inactivated FBS. The media were collected later for analysis by ELISA (as described below) for FVIII light chain antigen levels, and by the ChromZ FVIII as coagulation activity (FVIII-c) assay (Helena Labs) for biological activity, using the manufacturer's instructions and as described in Example 6.

10

15

EXAMPLE 5

Single Chain Factor VIII Infectivity Assay

20

25

30

In this Example, the infectivity of single chain Factor VIII was investigated. To determine the infectivity of rAAV-hF8-1, HepG2, 293, and H2.35 cells were infected with rAAV-hF8-1 and a control vector rAAV-hF8L at an MOI of 1×10^4 viral particles per cell. Recombinant AAV DNA in infected cells was isolated by Hirt extraction and run on an alkaline agarose gel. Southern blot analysis using an human F8 probe showed that similar amounts of rAAV-hF8-1 and rAAV-hF8L were isolated from uncoated virus in the infected cells. An infectious center assay (ICA) known in the art (*See e.g.*, Snyder, *Current Protocols in Genetics*, Chapter 12, John Wiley & Sons [1997]) was used to further characterize the infectivity of rAAV-hF8-1. In this assay, the infectious particle to total particle ratio of rAAV-F8-1 and that of a control

rAAV vector with the genome size of 4645 nucleotides was determined. The results indicated that rAAV-hF8-1 had an infectious particle to total particle ratio that was comparable to the control vector at approximately 1:1000. Taken together, these results indicate rAAV-hF8-1 has similar infectivity as rAAV vectors with the genome size of wild-type AAV.

EXAMPLE 6

Factor VIII Protein Expression Assay

An ELISA specific for the light chain of FVIII was used to determine FVIII light chain antigen levels in the 293 cells, as well as the injected animals (described below). NUNC Maxisorb 96 well plates were coated with 50 μ l of a 1:500 dilution of the light chain specific antibody, N77110 (Biodesign International) in a coating buffer overnight at 4°C. The plate was washed three times with wash buffer (PBS, 0.05% Tween 20) and blocked with 200 μ l blocking buffer (PBS, 10% horse serum, 0.05% Tween 20) at room temperature for 1 hour. The plate was washed three times and standards and samples were applied. Bioclone recombinant human FVIII (Baxter Hyland) was used as the standard, and was diluted in blocking buffer to concentrations ranging from 320 ng/ml to 10 ng/ml.

For analysis of transduced culture supernatants, the standards contained 50% media, and for analysis of mouse plasma, the standards were diluted into 10%, in normal pooled mouse plasma (Sigma). A standard assay reference plasma (SARP; Helena Labs) was also included in the assay. Following the loading of the standards and samples (95 μ l/well), the plate was incubated at room temperature for 2 hours, and washed five times with wash buffer (200 μ l/well). A 1:200 dilution of a horseradish peroxidase-conjugated light chain specific antibody, ESH8-HRP, (American Diagnostica) was added (100 μ l/well), and the plate was incubated for 1 hour at room temperature. The plates were then washed four times with wash buffer, and the antigen was detected using an ABTS peroxidase substrate kit (BioRad) according to the manufacturer's instructions. The results are shown in Table 1 of Example 7, below.

EXAMPLE 7

Factor VIII Biological Activity Assay

The ChromZ FVIII coagulation activity (FVIII-c) assay (Helena Labs, Beaumont, TX) was used to detect biologically active FVIII in the 293 cells infected as described in Example 4. Bioclone recombinant human FVIII (Baxter Hyland) was used as a standard to analyze transfected culture supernatants. The standards were diluted in plasma dilution buffer (supplied in kit) in the range of 10 ng/ml to 0.313 ng/ml, and were made 2.5% in media. Because this assay can detect both human and murine FVIII activity, it was modified to deplete biologically active human Factor VIII in the mouse plasma. Mouse plasma was pre-incubated with an antibody specific for human FVIII prior to performing the assay. The difference in FVIII activity between the untreated plasma sample and the antibody treated sample represent the amount of biologically active human FVIII in the plasma. The standard used in the assay was normal pooled human plasma (FACT; obtained from George King Biomedical). Serial dilutions of FACT were made in FVIII deficient plasma from undiluted (200 ng/ml) to 6.25 ng/ml. The standards (10 μ l) were incubated at 37°C for 15 min., with or without the addition of 2 μ l antibody N77110. Similarly, mouse plasma samples were diluted in FVIII deficient plasma and 10 μ l of these diluted samples were incubated with or without 2 μ l of N77110 at 37°C for 15 min., and immediately placed on ice. Thus, all incubations with antibody were done in a background of 100% plasma. The antibody adsorbed and non-adsorbed FACT standards, as well as the mouse plasma samples were diluted 1:20 in plasma detection buffer provided in the ChromZ kit. Thus, the final concentration of the FACT standards used in the assay ranged from 10 ng/ml to 0.313 ng/ml.

Twenty five microliters of these dilutions were added to a chilled 96 well plate. With the plate on ice, 25 μ l of FIXa reagent and 50 μ l of FX were added, and the plate was incubated at 37°C for 15 min. Substrate (50 μ l) was added and the plate was incubated for an additional 3 min at 37°C. The reaction was stopped with the addition of 25 μ l 50% acetic acid and the optical density at 405 nm was measured.

As shown below in Table 1, infection of 293 cells with AAV-e δ D-FVIII-HC resulted in no antigen production, as well as no biologically-active protein. Cells infected with AAV-e δ D-FVIII-LC produced FVIII light chain, but no biologically active protein. However, cells transduced with both vectors produced FVIII light chain and biologically active FVIII in a dose-dependent manner. Transduction of cells with the negative control vector, AAV-e δ D-FIX, resulted in no antigen nor any biologically active FVIII. It was assumed that equal amounts of heavy and light chains were produced in transduced cells. The activity units were converted to nanograms using the definition of one unit being equal to the amount of FVIII in 1 ml of normal pooled human plasma, or 200 ng.

**TABLE 1. *In Vitro* Production of Biologically Active
Human Factor VIII From Two rAAV Vectors**

	Vector	MOI	ELISA (ng/ml)	ChromZ	
				(mU/ml)	(ng/ml)
5	AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC	3×10^3	24	35	7.1
	AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC	3×10^4	121	440	87.9
	AAV-e δ D-FVIII-HC	3×10^3	0	0	0
10	AAV-e δ D-FVIII-HC	3×10^4	0	0	0
	AAV-e δ D-FVIII-LC	3×10^3	20.5	0	0
	AAV-e δ D-FVIII-LC	3×10^4	96.9	0	0
	AAV-hFIX	3×10^3	0	0	0
	AAV-hFIX	3×10^4	0	0	0
15	No Vector		0	0	0

EXAMPLE 8

Immunofluorescent Staining of FVIII Heavy and Light Chains

20 In these experiments, 293 cells transduced as described above were analyzed using immunofluorescent staining. 293 cells were plated on rat tail collagen-coated two-well culture slides at a density of 4×10^5 cells per well. Forty-eight hours later, the cells were transduced at an MOI of 3×10^4 particles per cell of rAAV-hFVIII-HC and rAAV-hFVIII-LC. Forty-eight hours post-transduction, the cells were fixed *in situ*

25 with acetone, blocked with 2% BSA, and stained with a fluorescently labelled anti-hFVIII light chain antibody and a fluorescently labelled anti-hFVIII heavy chain antibody. The anti-hFVIII light chain antibody used was ESH-4 monoclonal antibody (American Diagnostica), fluorescently labelled with alexa-488 (Molecular Probes),

according to the manufacturer's instructions. The anti-hFVIII heavy chain antibody used was MAS530P monoclonal antibody (Accurate Chemical) fluorescently labelled with alexa-594 (Molecular Probes), according to the manufacturer's instructions. The cells were counter-stained with DAPI. The images were collected using a Zeiss
5 Axioskop fluorescence microscope equipped with separate filters for DAPI, FITC, and rhodamine signals and a CCD camera. Image analysis was performed using Quips imaging software (Vysis).

As indicated above, infection of cells with either rAAV-e8D-FVIII-HC or AAV-e8D-FVIII-LC, followed by staining with antibodies to both chains resulted in
10 production of the individual chains of human FVIII. Immunofluorescent staining of cells co-infected with both vectors demonstrated that although some cells express only the heavy or light chain of hFVIII, many co-expressed both chains of human FVIII.

EXAMPLE 9

15 *In Vitro* Expression of Factor VIII Using Single Construct

Table 2 shows that two single vector constructs containing the heavy and the light chain of Factor VIII driven by different promoters express biologically active Factor VIII. The constructs pAAV-hF8-1 (SEQ ID NO:13), and pVm4.1cF8ΔB (SEQ ID NO:14) were transfected into 293 cells. Following transfection, the cells were
20 allowed to express factor VIII for 48-72 hours. Factor VIII in the culture media was assayed by the ChromZ FVIII coagulation activity (FVIII-c) assay, as per the manufacturer's instructions.

TABLE 2

In Vitro Production of Biologically Active FVIII

Construct(s)	ELISA (ng/ml)	ChromZ (ng/ml)
Control	-	0
pAAV-hF8-1	-	4.9
pVm4.1cF8ΔB	-	46

EXAMPLE 10

Factor VIII Expression Using Tissue Specific Promoters

In these experiments, different promoters and enhancer elements were used to drive expression of a Factor VII coding sequence. Expression of Factor VIII was compared in 293 cells and HepG2 cells using different promoters. The pAAVeF8ΔB contains an EF-1α promoter with a hGH intron, Factor VIII with a B-domain deletion (F8ΔB) and a polyA. As described previously, pAAV-hF8-1 uses the HNF-3 albumin promoter with a minimal intron followed by F8ΔB and a minimal polyA. The construct pAAV-c8 uses the CMV enhancer-promoter and the F8ΔB. pAAV8b1 contains the HNF-3 albumin promoter followed by the CMV/B-globin intron with the F8ΔB and a minimal poly A site. Table 3 describes Factor VIII expression using the albumin promoter relative to the control plasmid pV4.1cF8ΔB in HepG2 and 293 cells. These data show increased expression of Factor VIII in HepG2 liver cells with the albumin promoter as compared to Factor VIII expression in 293 cells.

TABLE 3

Relative Tissue Specificity of Promoters

Plasmid Construct	HepG2 Cells	293 Cells
pAAV-hF8-1	6.2	0.6
pAAV8bl	6.7	1.0
pAAVc8	30.0	41.0
pV4.1eF8ΔB	100	100

Next, several promoters derived from the transthyretin (TTR) gene promoter were transfected into HepG2 cells. TTR is an abundant serum protein and the gene enhancer-promoter contains well known liver-specific transcription factor binding sites (Samadani *et al.*, Gene Expression 6:23 [1996]; Yan *et al.*, EMBO 9:869 [1990]; Costa and Grayson, Nuc. Acids Res., 19:4139 [1991]; Costa *et al.*, Mol. Cell. Bio., 6:4697 [1986]). The constructs were made by replacing the HNF-3 albumin promoter in pAAV-hF8-1 with various lengths of the TTR promoter-enhancer. The TTR enhancer-promoter was modified by replacing the weak affinity binding sites with the strong affinity binding sites to create pAAV-hF8-2. The pAAV-hF8-TTR-E-L-P202 construct contains the full TTR promoter with a linker between the enhancer and the promoter. The remaining constructs are 5' deletions: pAAV-hF8-TTR-E-P202 has the promoter and enhancer with no linker; pAAV-hF8-TTR-E-P197 has a 5 base pair deletion from the promoter; pAAV-hF8-TTR-E-P151 has a 50 base pair deletion; pAAV-hF8-TTR-P202 lacks the TTR enhancer and pAAV-hF8-TTR(X) has a 65 base pair deletion in the enhancer. The control plasmid, pAAV-hF8-1, expressed approximately 4.6 mU/ml. Table 4 shows the fold-increase in Factor VIII activity using the TTR promoter series relative to the control plasmid.

TABLE 4

Factor VIII Expression Using TTR-Derived Promoters

Plasmid Construct	Relative Factor VIII Activity
pAAV-hF8-STTR	3.16
pAAV-hF8-TTR-E-L-P202	8.86
pAAV-hF8-TTR-E-P202	6.1
pAAV-hF8-TTR-EP197	7.3
pAAV-hF8-TTR-E-P151	13.3
pAAV-hF8-TTR-P202	2.3

EXAMPLE 11

In Vivo Expression of Factor VIII

In order to test the feasibility of the AAV vector approach of the present invention *in vivo*, three groups of five C57BL/6 mice were injected via the portal vein with either 3×10^{11} particles of AAV-e δ D-FVIII-HC, 3×10^{11} particles of AAV-e δ D-FVIII-LC, or 3×10^{11} particles of both AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC. In addition, a group of four animals was injected with 3×10^{11} particles of AAV-e δ D-FLX. It has been shown that this strain of mice does not elicit an immune response to human FVIII when the gene is delivered to the portal vein via an adenoviral vector (Connelly *et al.*, Blood 87:4671-4677 [1996]). As indicated by the results shown below, the data obtained during these experiments demonstrate the feasibility of producing biologically active FVIII using two AAV vectors to independently deliver the heavy and light chains of FVIII.

Blood samples were collected in sodium citrate via the retro-orbital plexus at biweekly intervals for the first 2 months and at monthly intervals thereafter for 6

months and at 11 months. Very high levels of FVIII light chain were expressed in animals injected with AAV-e δ D-FVIII-LC alone or both vectors as shown in Figure 8.

In order to assess the amount of biologically active human FVIII produced in the animals, a modified ChromZ assay was used. Since this assay detects both human and murine FVIII, the amount of FVIII present in the plasma before and after adsorption to an antibody specific to human FVIII was determined. The amount of FVIII remaining in the plasma after adsorption represented the amount of active murine FVIII and the difference represented the amount of active human FVIII. Control experiments demonstrated that the antibody could remove 80-90% of the human FVIII from a mouse plasma sample when the sample was spiked with up to 32 ng of human FVIII. The modified ChromZ assay indicated that only those animals injected with both vectors produced biologically active FVIII, as indicated in Table 5. The results shown in Table 5 are those from plasma collected 8 weeks post-injection, although similar results were obtained at 10 weeks and 5 months post-injection. One of the five animals co-injected with both the heavy and light chain vector did not express VIII, presumably due to an inefficient injection, and was omitted from the analysis. Animals injected with both vectors produced over 2 μ g/ml hFVIII light chain as measured by ELISA. The ChromZ assay indicated that a total of 600-900 ng/ml of active hFVIII was detected in the plasma. The contribution from murine Factor VIII was approximately 400-500 ng/ml, indicating that about 230-430 ng/ml of active human Factor VIII was present in the plasma. Although only a fraction of the total protein was found to be active, the animals produced physiological levels of the active protein (*i.e.*, 200 ng/ml). The animals were found to have maintained these physiological levels of active protein for more than 11 months, without waning. Similar analyses performed on animals injected with the light chain vector alone, the heavy chain vector alone, or the hFIX vector demonstrated no biologically active human FVIII in the plasma of these animals.

TABLE 5

Biological Activity of Human Factor VIII *In Vivo*

Construct(s) Used	ELISA (ng/ml)	Total FVIII (-Ab) (Units)	Murine FVIII (+Ab) (Units)	Human FVIII (ng/ml)
5 AAV-e δ D-FVIII-HC and AAV-e δ D-FVIII-LC*	2288	3.9	2.2	342
AAV-e δ D-FVIII-LC *	3329	1.4	1.6	0
AAV-e δ D-FVIII-HC	0	1.6	1.6	0
10 AAV-e δ D-FLX	0	1.4	2.0	0

*Average of three animals.

EXAMPLE 12

15 Gene Transfer and Vector Expression in Tissues

In these experiments, evidence of gene transfer to liver was obtained by Southern Blot analysis following isolation of DNA from one animal of each experimental group sacrificed 8 weeks post-injection (*i.e.*, as described in Example 11). In addition, DNA was obtained from other tissues in order to determine the
20 degree of vector expression in organs other than the liver.

Twenty micrograms of DNA was digested with *Bgl*II, separated using a 1% agarose gel, and hybridized with a ³²P-labelled 1126 bp *Al*wNI fragment encoding the A1 and A2 domains of hFVIII (heavy chain probe), or a ³²P-labelled 1456 bp *Nde*I-*Eco*RI fragment encoding the A3, C1 and C2 domains of hFVIII (light chain probe).
25 Copy number controls were generated by spiking *Bgl*II-digested naive mouse liver DNA with *Bgl*II-digested heavy or light chain plasmid DNA (pVm4.1e δ D-hFVIII-HC and pVm4.1e δ D-hFVIII-LC, respectively), at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome. The hybridized membranes were analyzed using a Storm 860 phosphorimager (Molecular Dynamics), and quantitation of vector copy number was
30 evaluated using ImageQuaNT software (Molecular Dynamics). Autoradiography of the

hybridized membranes was also performed. Total RNA was isolated from liver tissue using the RNA Stat extraction kit (Tel-Test). As describe briefly below, Northern blot analysis was also performed on 10 µg RNA using methods known in the art, in conjunction with the ³²P-labelled probes specific to the heavy and light chains of hFVIII described above and autoradiography was performed on the hybridized membranes.

Following digestion of liver DNA with *Bgl*II and hybridization with an hFVIII light chain probe described below, using methods known in the art, a band at the predicted size of 3015 bp was detected in animals injected with rAAV-hFVIII-LC, or both the heavy and light chain vectors. This band was not observed in the DNA of animals injected with the heavy chain vector alone or the hFIX vector, as shown in Figure 9A (rAAV-hFVIII-LC, lane 1; rAAV-hFIX, lane 2; rAAV-hFVIII-HC, lane 3; both rAAV-hFVIII-LC and rAAV-hFVIII-HC, lane 4; copy number controls were generated by spiking *Bgl*II digested naive mouse liver DNA with the corresponding plasmids at ratios of 10, 5, 1, 0.1, and 0.01 copies per diploid genome, lanes 5-9). Phosphoimage analysis revealed that the light chain vector was present at approximately 2.4 and 1.5 copies per diploid genome in animals injected with the light chain vector alone or both vectors, respectively. When *Bgl*II-digested DNA was hybridized with an hFVIII heavy chain probe, the expected band of 2318 bp was observed in animals injected with the heavy chain vector alone or both vectors, but was not detected in animals injected with the light chain vector alone or the hFIX vector, as shown in Figure 9B. The copy number in animals injected with the heavy chain vector alone and both vectors was 1.1 and 1.7 vector copies per diploid genome, respectively.

The results of hybridization of DNA extracted from the spleen, kidney and heart tissue with either an hFVIII light chain probe or a heavy chain probe indicated that these tissues contained less than 1 copy of vector sequences per 10 diploid genomes, demonstrating that the vector distributes primarily to the liver following intra-portal injection, as shown in Figures 10A and 10B.

Human FVIII gene expression in the liver of the mice was also assessed by Northern blot analysis on RNA isolated from animals sacrificed 8 weeks post-injection (as described above). hFVIII light chain transcripts of the predicted size (2.7 kb) were observed in animals injected with the light chain vector alone or both vectors, as
5 shown in Figure 11A. Similarly, the expected hFVIII heavy chain transcripts (2.7 kb) were detected in animals that were injected with the heavy chain vector alone or both vectors, as shown in Figure 11B. Since the heavy and light chain DNA sequences were shown by Southern blot analysis to be present at approximately the same copy number (1.7 and 1.5 copies per diploid genome, respectively), in an animal injected
10 with both vectors, these results demonstrate that both the heavy and light chains of hFVIII are expressed in the liver in approximately equivalent amounts.

CLAIMS

We claim:

1. A method of treating hemophilia in a human comprising:
 - 5 a) providing at least one recombinant adeno-associated vector comprising nucleotide sequences encoding Factor VIII; and
 - b) administering said at least one recombinant adeno-associated vector into said subject under conditions such that said Factor VIII nucleotide sequences are expressed at a level which
 - 10 provides a therapeutic effective amount in said subject.
2. The method of Claim 1, wherein said Factor VIII nucleotide sequences are expressed in the liver of said subject.
- 15 3. The method of Claim 1, wherein said blood clotting disorder is ameliorated.
4. The method of Claim 1, wherein said blood clotting disorder is hemophilia A.
- 20 5. The method of Claim 1, wherein said subject is a human.
6. The method of Claim 1, wherein said recombinant adeno-associated vector encodes the light chain of said Factor VIII.
- 25 7. The method of Claim 1, wherein said recombinant adeno-associated vector encodes the heavy chain of said Factor VIII.

8. The method of Claim 1, wherein said Factor VIII nucleotide sequences encode at least one Factor VIII domain selected from the group consisting of A1, A2, A3, B, C1, and C2.

5 9. The method of Claim 1, wherein said recombinant adeno-associated vector is administered intravenously.

10 10. The method of Claim 8, wherein said intravenous administration is via the portal vein.

11. The method of Claim 1, wherein said recombinant adeno-associated vector further comprises nucleotides encoding the human growth hormone polyadenylation sequence.

15 12. A method of treating a subject comprising:

a) providing:

i) a subject suffering from a blood clotting disorder,

ii) a first recombinant adeno-associated vector comprising nucleotide sequences encoding the light chain of Factor VIII, and

20

iii) a second recombinant adeno-associated vector comprising nucleotide sequences encoding the heavy chain of Factor VIII; and

b) administering said at first and second recombinant adeno-associated vector into said subject under conditions such that said Factor VIII heavy and light chain nucleotide sequences are expressed at a level which provides a therapeutic effect in said subject.

25

13. The method of Claim 12, wherein said Factor VIII heavy and light chain nucleotide sequences are expressed in the liver of said subject.

14. The method of Claim 12, wherein said Factor VIII nucleotide sequences
5 encode at least one Factor VIII domain selected from the group consisting of A1, A2, A3, B, C1, and C2.

15. The method of Claim 12, wherein said blood clotting disorder is ameliorated.

10

16. The method of Claim 12, wherein said blood clotting disorder is hemophilia A.

15

17. The method of Claim 12, wherein said subject is a human.

18. The method of Claim 12, wherein said recombinant adeno-associated vector is administered intravenously.

19. The method of Claim 18, wherein said intravenous administration is via
20 the portal vein.

25

20. The method of Claim 12, wherein said recombinant adeno-associated vector further comprises nucleotides encoding the human growth hormone polyadenylation sequence.

21. A recombinant adenovirus-associated vector, wherein said vector comprises nucleic acid encoding at least a portion of Factor VIII operably linked to

control sequences, wherein said control sequences direct the transcription of translation of said portion of Factor VIII, and a pharmaceutically acceptable excipient.

22. The vector of Claim 21, wherein said nucleic acid encoding Factor VIII
5 comprises nucleotides encoding the heavy chain of Factor VIII.

23. The vector of Claim 21, wherein said nucleic acid encoding Factor VIII
comprises nucleotides encoding the light chain of Factor VIII.

10 24. The vector of Claim 21, wherein said nucleic acid encoding at least a
portion of Factor VIII encodes at least one Factor VIII domain selected from the group
consisting of A1, A2, A3, B, C1, and C2.

25. The vector of Claim 21, wherein said vector further comprises
15 nucleotides encoding the human growth hormone polyadenylation sequence.

26. A host cell containing at least one recombinant adeno-associated vector,
wherein said host cell is capable of expressing Factor VIII.

20 27. The host cell of Claim 26, wherein said cell is a liver cell.

28. The host cell of Claim 26, wherein said Factor VIII is expressed at or
above physiological levels.

25 29. The host cell of Claim 26, wherein said host cell expresses said Factor
VIII at a level greater than about 200 ng/ml.

30. The host cell of Claim 29, wherein said host cell is present in an animal.

FIGURE 1




FIGURE 2





FIGURE 3

AAV-F8-1



BEST AVAILABLE COPY

FIGURE 4

PVM4.1c-F8AB




FIGURE 5

CAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCGGGCGTGGGGCGACCTTTGGTGGCCCGGCCTCAGT
GAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGGCCCGCCAGGGAATGTTTGTCTT
AAATACCATCCAGGGAATGTTTGTCTTAAATACCATCCAGGGAATGTTTGTCTTAAATACCATCTACAGTTATTGGTT
AAAGAAGTATATTAGAGCGAGTCTTTCTGCACACAGATCACCTTTCCGGGTGCCGCCCTTAGGCAGGTAAAGTGCCGTGTG
TGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCCTTGGCTGCCCTTGAATTACTGACACTGACATCCACTTTTTCT
TTTTCTCCACAGGTATCGATTCCACCATGCAAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCCTTTTGGGATTCTGCTTT
AGTGCCACCAGAAGATACTACCTGGGTGCAGTGGAACTGTGATGGGACTATATGCAAAGTGATCTCGGTGAGCTGCCTGT
GGACGCAAGATTTCTCTCTAGAGTGCCAAAATCTTTCCATTCAACACCTCAGTCGTGTACAAAAGACTCTGTTTGTAG
AATTACGGATCACCTTTTCAACATCGCTAAGCCAAGGCCACCTGGATGGGTCTGCTAGGTCTTACCATCCAGGCTGAG
GTTTATGATACAGTGGTCATTACACTTAAGAACATGGCTTCCATCCTGTCAGTCTTCATGCTGTTGGTGTATCCTACTG
GAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAAAGAAGATGATAAAGTCTTCCCTGGTGGAA
GCCATACATATGTCTGGCAGGTCTGAAAGAGAATGGTCCAATGGCCTCTGACCCACTGTGCCCTTACCTACTCATATCTT
TCTCATGTGGACCTGGTAAAGACTTGAATTACGGCCTCATTGGAGCCCTACTAGTATGTAGAGAAGGAGTCTGGCCAA
GGAAAGACACAGACCTTGCAAAAATTTATACTACTTTTGTCTGTATTGATGAAGGGAAAAGTTGGCACTCAGAAACAA
AGAACTCCTTGATGTCAGGATAGGGATGCTGCATCTGCTCGGGCCTGGCCTAAAATGCACACAGTCAATGGTTATGTAAC
AGGTCTCTGCCAGGTCTGATTGGATGCCACAGGAATCAGTCTATTGGCATGTGATTGGAATGGGCACCACTCCTGAAGT
GCACCTCAATTTCTCGAAGGTGCACACTTTCTGTGAGGAACCATCGCCAGGCGTCTCGCAAAATCTGCCAATAACTT
TCCTTACTGCTCAAACACTCTTGATGGACCTTGGACAGTCTTCTACTGTTTGTGTCATATCTTCCCAACATCATGTGGC
ATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCCCACTACGAATGAAAAATAATGAAGAAGCGGAAGACTA
TGATGATGATCTTACTGATTCTGAAATGGATGTGGTCAGGTTTGTATGATGACAACCTCTCCTTCTTTATCCAAATTCGCT
CAGTTGCCAAGAAGCATCCTAAAACCTGGGTACATTACATTGCTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTC
CTCGCCCCGATGACAGAAGTTATAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAGGAAGTACAAAAAGT
CCGATTATGGCATAACAGATGAAACCTTTAAGACTCGTGAAGCTATTACGATGAATCAGGAATCTTGGGACCTTTAC
TTTATGGGGAAGTTGGAGACACACTGTGTGATTATTTAAGAATCAAGCAAGCAGACCATATAACATCTACCTCAGCGGA
ATCACTGATGTCCGTCCTTTGTATTCAAGGAGATTACCAAAAGGTGTAACAACTTTGAAGGATTTTCCAAATCTGCCAGG
AGAAATATTCAAATATAAATGGACAGTGAAGTGTAGAAGATGGGCCAACTAAATCAGATCCTCGGTGCTGACCCGCTATT
ACTCTAGTTTCTGTTAATATGGAGAGAGATCTAGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATCTGTA
GATCAAAGAGGAAACAGATAATGTGAGACAAGAGGAATGTGATCCTGTTTCTGTATTGATGAGAACCGAAGCTGGTA
CCTCACAGAGAATATACAACGCTTTCTCCCAATCCAGCTGGAGTGCAGCTTGAGGATCCAGAGTTCCAAGCCTCCAACA
TCATGCACAGCATCAATGGCTATGTTTGTATAGTTTGCAGTTGTGAGTTTGTGTCATGAGGTGGCATACTGGTACATT
CTAAGCATTTGGAGCACAGACTGACTTCTTCTGCTCTTCTTCTCTGGATATACCTTCAAACACAAAATGGTCTATGAAGA
CACACTCACCTTATTCCCATTTCTCAGGAGAACTGTCTTCATGTGATGGAACCCAGGTCTATGGATTCTGGGGTGCC
ACAACCTCAGACTTTCCGGAACAGAGGCATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACACTGGTGATTATTAC
GAGGACAGTTATGAAGATATTTACGATACTTGTCTGAGTAAAAACAATGCCATTGAACCAAGAAGCTTCGAAATAACTCG
TACTACTCTTCAGTCAGATCAAGAGGAAATTGACTATGATGATACCATATCAGTTGAATGAAGAAGGAGATTTTGACA
TTTATGATGAGGATGAAATCAGAGCCCCCGCAGCTTTCAAAGAAAAACAGACACTATTTTATGCTGTCAGTGGAGAGG
CTCTGGGATTATGGGATGAGTAGCTCCCAATGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAA
AGTTGTTTTCAGGAATTTACTGATGGCTCCTTTACTCAGCCCTTATACCGTGGAGAACTAAATGAACATTTGGGACTCC
TGGGGCCATATATAAGAGCAGAAGTTGAAGATAATATCATGGTAACCTTTCAGAAATCAGGCCTCTCGTCCCTATTCTTC
TATTCTAGCCTTATTTCTATGAGGAAGATCAGAGGCAAGGAGCAGAACCTAGAAAAAACTTTGTCAAGCCTAATGAAAC
CAAACTTACTTTTGGAAAGTGCAACATCATATGGCACCCTAAAGATGAGTTTGACTGCAAAGCCTGGGCTTATTTCT
CTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCTGGTCTGCCACACTAACACACTGAACCT
GCTCATGGGAGACAAGTGACAGTACAGGAATTTGCTCTGTTTTCACCATCTTTGATGAGACCAAAAGCTGGTACTTCAC
TGAAAAATATGGAAGAACTGCGAGGCTCCCTGCAATATCCAGATGGAAGATCCCACTTTTAAAGAGAATTATCGCTTCC
ATGCAATCAATGGCTACATAATGGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTCGATGGTATCTGCTC
AGCATGGGCAGCAATGAAAAATCCATTCTATTCTAGTGGACATGTGTTCACTGTACGAAAAAAGAGGAGTATAA
AATGGCACTGTACAATCTCTATCCAGGTGTTTTGAGACAGTGGAAATGTTACCATCCAAAGCTGGAATTTGGCGGGTGG
AATGCCTTATGGCGAGCATCTACATGCTGGGATGAGCACACTTTTTCTGGTGTACAGCAATAAGTGTGAGACTCCCTG
GGAATGGCTTCTGGACACATTAGAGATTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCAAAGCTGGCCAG
ACTTCATTATTCGGATCAATCAATGCTGGAGCACCAGGAGCCCTTTCTTGGATCAAGGTGGATCTGTTGGCACCA
TGATTATTACGGCATCAAGACCCAGGGTGCCCGTCAGAAGTTCTCCAGCCTCTACATCTCTCAGTTTATCATCATGTAT
AGTCTTGATGGGAAGAAGTGGCAGACTTATCGAGGAAATTCACCTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATT
ATCTGGGATAAAACACAATATTTTAACCTCCAATTATTGCTCGATACATCCGTTTGACCCCACTCATTATAGCATTC

GCAGCACTCTTCGCATGGAGTTGATGGGCTGTGATTTAAATAGTTGCAGCATGCCATTGGGAATGGAGAGTAAAGCAATA
TCAGATGCACAGATTACTGCTTCATCCTACTTTACCAATATGTTTGCCACCTGGTCTCCTTCAAAGCTCGACTTCACCT
CCAAGGGAGGAGTAATGCCTGGAGACCTCAGGTGAATAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGA
AAGTCACAGGAGTAATACTACTCAGGGAGTAAAATCTCTGCTTACCAGCATGTATGTGAAGGAGTTCTCATCTCCAGCAGT
CAAGATGGCCATCAGTGGACTCTCTTTTTTTCAGAATGGCAAAGTAAAGGTTTTTTCAGGGAAATCAAGACTCCTTCACACC
TGTGGTGAACCTCTCTAGACCCACCGTTACTGACTCGCTACCTTCGAATTCACCCCAAGAGTTGGGTGCACCAGATTGCCC
TGAGGATGGAGGTTCTGGGCTGCGAGGCACAGACCTCTACTGACTCGAGAATAAAAGATCAGAGCTCTAGAGATCTGTG
TGTTGGTTTTTTGTGTGCGGCCGAGGAACCCGATTAGTAGAGGTTGGCCACTCCCTCTCTGCGCGCTCGCTCGCTCACT
GAGGCCGGGCGACCAAAGGTCGCCCCGACGCCCGGGCTTTGCCCGGGCGGCTCAGTGAGCGAGCGAGCGCGCAGCTGCCT
GCAGGACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAGGCCGCTTGTGCGGTTTTTTCATAGGCTCC
GCCCCCTGACGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCG
TTTCCCCCTGGAAGCTCCCTCGTGCCTCTCTGTTCCGACCTGCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTC
GGGAAGCGTGGCGCTTTCTCATAGCTCAGCTGTAGGTATCTCAGTTCCGTGTAGGTCTGCTCCAAGCTGGGCTGTG
TGCACGAACCCCCGTTCCAGCCGACCGCTGCGCTTATCCGGTAACATATCGTCTTGAGTCCAAACCCGGTAAGACACGAC
TTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGGTATGTAGGCGGTCTACAGAGTTCTTGAAGTG
GTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAG
TTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGTGTTGCAAGCAGCAGATTACGCGCAGA
AAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGCTCTGACGCTCAGTGGAACGAAAACCTCACGTTAAGGGAT
TTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAAATGAAGTTTTAAATCAATCTAAAGTA
TATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTCGTTCA
TCCATAGTTGCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCGCAGTGCTCGAATGAT
ACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTC
CTGCAACTTTATCCGCTCCATCCAGTCTATTAATTTGTTGCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTG
CGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCGTCTGTTGGTATGGCTTCATTCAGCTCCGGTTCCCA
ACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAA
GTAAGTTGGCCGAGTGTATCACTCATGTTATGGCAGCACTGCATAATTCTTACTGTCTATGCCATCCGTAAGATGC
TTTTCTGTGACTGGTGAAGTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGCGCTC
AATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACCTC
CAAGGATCTTACCGCTGTTGAGATCCAGTTGATGTAACCCACTCGTGACCCCACTGATCTTCAGCATCTTTTACTTTT
ACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAAGGGCGACAGGAAATGTTGAAT
ACTCATACTCTTCTTTTTCAATATTATGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTA
TTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTTATTATC
ATGACATTAACCTATAAAAATAGGCGTATCACGAGGCCCTTTCTGCTCGCGCGTTTCGGTGATGACGGTGAACCTCTG
ACACATGCAGCTCCCGGAGACGGTACAGCTTGTCTGTAAGCGGATGCCGGGAGCAGACAGCCCGTCAGGGCGCGCTCAG
CGGGTGTGCGGGGTGTCGGGGCTGGCTTAACATGCGGCATCAGAGCAGATTGTACTGAGAGTGACCATATAAATTGTA
AACGTTAATATTTTGTAAAAATTGCGGTTAAATTTTTGTTAAATCAGCTCATTTTTTTAACCAATAGGCCGAAATCGGCAA
AATCCCTTATAAATCAAAAGATAGCCCGAGATAGGGTTGAGTGTGTTCCAGTTTGGAAACAAGAGTCCACTATTAAAGA
ACGTGGACTCCAACGTCAAAGGGCGAAAACCGTCTATCAGGGCGATGCGCCACTACGTGAACCATCACCCAAATCAAGT
TTTTTGGGGTCCAGGTGCCGTAAAGCACTAAATCGGAACCTAAAGGGAGCCCCGATTAGAGCTTGACGGGGAAGGCC
GGCGAACGTGGCGAGAAAGGAAGGAAGGAAGCGAAAGGAGCGGGCGCTAGGGCGCTGGCAAGTGTAGCGGTCAACGCTGC
GCGTAACCAACACACCCCGCGCTTAATGCGCCGCTACAGGGCGGCTACTATGGTTGCTTTGACGTATGCGGTGTGAAA
TACCGCACAGATGCGTAAGGAGAAAAATACCGCATCAGGCCGTAACCTGTCCGATCACCGGAAAGGACCCGTAAAGTGATA
ATGATTATCATCTACATATCACACGTGCGTGGAGGCCATCAAACCACGTCAAATAATCAATTATGACGCAGGTATCGTA
TTAATTGATCTGCATCAACTTAACGTAAAAACAACCTCAGACAATACAAATCAGCGACACTGAATACGGGGCAACCTCAT
GTCAACGAAGAACAGAACCCCGAGAACAAACCCGCAACATCCGCTTTCTTAACCAATGATTGAACAAATTAACATCG
CTCTTGAGCAAAAAGGGTCCGGGAATTTCTCAGCCTGGTTCATTGAAGCCTGCCGTCCGAGACTAACGTCAGAAAAGAGA
GCATATACATCAATTAAAGTGATGAAGAATGAACATCCCGGCTTCTCCCTCCGAACAGGACGATATTGTAATTCACCT
TAATTACGAGGGCATTGCAATTAATTGAGTTGAGTTTACCCTTTCTGACAGTGACAGACTGCGGTGTTGGCTCTGTCA
CAGACTAAATAGTTGAATGATTAGCAGTTATGGTGATCAGTCAACCACCAGGGAATAATCCTTCATATTATTATCGTGC
TTCACCAACGCTGCTCAATTGCTCTGAATGCTTCCAGAGACACCTTATGTTCTATACATGCAATTACAACATCAGGGTA
ACTCATAGAAATGCTGCTATTAGCATATTTTTTACACGAATCAGATCCACGGAGGGATCATCAGCAGATTGTTCTTTAT
TCATTTTGTGCTCCATGCGCTTGTCTTCTCATCTAGCGGTTAAATATTACTTCAAATCTTTCTGTATGAAGATTTGAGC
ACGTTGGCCCTTACATACATCTGTGCGTTGTATTTCCCTCAGAAATGCCAGCAGGACCGCACTTTGTTACGCAACCAATAC
TATTAAGTGAAAACATTCCTAATATTGACATAAATCATCAACAAAACACAAGGAGGTGAGCCAGATTGAACGATAAA
AACGATAATGCAAACTACGCGCCCTCGTATCACATGGAAGTTTTTACCAATGGCTCAGGTTGCCATTTTTTAAAGAAATAT
TCGATCAAGTGCAGAAAGATTTAGACTGTGAATGTTTTTATTCTGAACATAAACGCTCACACGCTCTCACATTATATTAC
TATCTAGCCACAGATAATATTACATCGTGTAGAAAACGATAACACCGTGTAAATAAAGGACTTAAAAAGGTTGTAA

TGTTAAATTCTCAAGAAACACGCATCTTATAGAAACGTCCTATGATAGGTTGAAATCAAGAGAAATCACATTTTCAGCAAT
ACAGGGGAAAATCTTGCTAAAGCAGGAGTTTTCCGATGGGTTACAAATATCCATGAACATAAAAGATATTACTATACCTTT
GATAATTCATTACTATTTACTGAGAGCATTACAGAACTACACAAATCTTTCCACGCTAAATCATAACGTCGGGTTTCTT
CCGTGTCAGCACCGGGCGTGGCATAATGCAATACGTGTACGCGCTAAACCCTGTGTGCATCGTTTTAATTATTTCCCGG
ACACTCCCGCAGAGAAGTTCCCGTCAGGGCTGTGGACATAGTTAATCCGGGAATACAATGACGATTCATCGCACCTGAC
ATACATTAAATAATATTAACAAATATGAAATTTCAACTCATTGTTTAGGGTTTGTTTAATTTTCTACACATACGATTCTGC
GAACTTCAAAAAGCATCGGGAATAACACCATGAAAAAATGCTACTCGCTACTGCGCTGGCCCTGCTTATTACAGGATGT
GCTCAACAGACGTTTACTGTTCAAAACAAACCGGCAGCAGTAGCACCAAAGGAAACCATCACCCATCATTCTTCGTTTC
TGGAATTGGGCAGAGAAAACGTGTCGATGCAGCCAAATTTGTGGCGGCGCAGAAAAATGTTGTTAAACAGAAACCCAGC
AAACATTTCGTAAATGGATTGCTCGGTTTTATTACTTTAGGCATTTATACTCCGCTGGAAAGCGCGTGTGTATTGCTCACA
TAATTGCATGAGTTGCCATCGCATATGGGCAACTCTATCTGCACTGCTCATTAAATATACTTCTGGGTTCTTCCAGTT
GTTTTTGCATAGTATCAGCCTCTCTCTGAGGGTGAAATAATCCCGTTTCAGCGGTGTCTGCCAGTCGGGGGAGGCTGCA
TTATCCACGCGCGAGGCGGTGTGGCTTCACGCACTGACTGACAGACTGCTTTGATGTGCAACCGACGACGACCAGCGGC
AACATCATCACGCGAGCATCATTTTCAGCTTTAGCATCAGCTAACTCCTTCGTGTATTTTGCATCGAGCGCAGCAACAT
CAGGCTGACGCATCTGCATGTCAGTAATTGCCCGGTTCCGCCAGCTTCAGTTCTCTGGCATTTTTGTGCGCTGGGCTTTG
TAGGTAATGGCGGTATCACGGTAATGATTAAACAGCCCATGACAGGCGAGCATGATGCAGATAACCAGAGCGGAGATAAT
CGCGGTGACTCTGCTCATACATCAATCTCTGACCGTTCCGCCCGCTTCTTTGAATTTTGAATCAGGCTGTCTAGCCTT
ATGCTCGAACTGACCATAACCAGCGCCCGGCGAGTGAAGCCAGATATTGCTGCAACGGTCGATTGCTGACGGATATCAC
CAGCATCAATCATAGGTAAAGCGCCACGCTCCTTAATCTGCTGCAATGCCACAGCGTCCTGACTTTTCGGAGAGAAGTCT
TTCAGGCCAAGCTGCTTGGCGTAGGCATCCCAACCGGAAAGAGCTGGTAGCGTCCGGCGCCTGTTGATTTGAGTTT
TGGGTTTAGCGTGACAAGTTTGGCAGGGTGATCGGAGTAATCAGTAAATAGCTCTCCGCCTACAATGACGTCTAACCAT
GATTTCTGGTTTTCTGACGTCCGTTATCAGTTCCCTCCGACCAGCCAGCATATCGAGGAACGCCCTTACGTTGATTATTG
ATTTCTACCATCTTCTACTCCGCTTTTTTAGCAGCGAAGCGTTTGATAAGCGAACCATCGAGTCAGTACCGATGTAGC
CGATAAACACGCTCGTTATATAAGCGAGATTGCTACTTAGTCCGGCGAAGTCGAGAAGGTCACGAATGAACCAGGCGATA
ATGGCGCACATCGTTGCGTCGATTACTGTTTTTGTAAACGACCGCCATTATATCTGCCGCGAAGGTACGCCATTGCAAA
CGCAAGGATTGCCCCGATGCCTGTTCCTTTGCCGCGAGAATGGCGGCCAACAGGTCATGTTTTTCTGGCATCTTCATGT
CTTACCCCCAATAAGGGGATTGCTCTATTTAATTAGGAATAAGGTCGATTACTGATAGAACAAATCCAGGCTACTGTGT
TTAGTAATCAGATTTGTTCTGACCGATATGCACGGGCAAAACGGCAGGAGTTGTTAGCGCGACCTCCTGCCACCCGCT
TTCAGGAAGGTCATGTGTAAAGGCCGCGAGCGTAACTATTACTAATGAATTCAGGACAGACAGTGGCTACGGCTCAGTTT
GGGTTGTGCTGTGTGCTGGGCGCGATGACGCGCTGTACGCATTGGTGATCCGGTTCTGCTTCCGGTATTGCTTAATTCA
GCACAACGGAAGAGCACTGGCTAACAGGCTCGCCGACTCTTCAGGATTATCGACTCAATGCTCTTACCTGTTGTGCAG
ATATAAAAAATCCGAAACCGTTATGCAGGCTCTAACTATTACCTGCGAACTGTTTCGGGATTGCATTTTGACAGCTCT
CTGCCTGCGATGGTTGGAGTTCAGACGATACGTGGAAGTGACCAACTAGGCGGAATCGGTAGTAAGCGCCGCTCTTTT
CATCTCACTACCACAACGAGCGAATTAACCCATCGTTGAGTCAAAATTTACCCAAATTTATTCAATAAGTCAATATCATGC
CGTTAATATGTTGCCATCCGTGGCAATCATGCTGCTAACGTGTGACCGCATTCAAAATGTTGTCTGCGATTGACTCTTCT
TTGTGGCATTGCACCACCAGAGCGTCATACAGCGGCTTAACAGTGCGTGACCAGGTGGGTTGGSTAAGGTTTGGGATTAG
CATCGTCACAGCGCGATATGCTGCGCTTGCTGGCATCCTTGAATAGCCGACGCTTTGTCATCTTCCGCACTCTTTCTCGA
CAACTCTCCCCACAGCTCTGTTTTGGCAATATCAACCGCACGGCCTGTACCATGGCAATCTCTGCATCTTGGCCCCGGC
GTCGCGGCACTACGGCAATAATCCGCATAAGCGAATGTTGCGAGCACTTGCAGTACCTTTGCCTTAGTATTTCTTCAAG
CTGCCCCCTGCAGG

FIGURE 6

CGCCCCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCGGGCAA
AGCCCGGGCGTCGGGCGACCTTTGGTCGCCCCGGCCTCAGTGAGCGAGCGAGC
GCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTGCGGCCGCACG
CGTGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCT
TTTTCCCAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAAC
GTTCTTTTTTCGCAACGGGTTTGCCGCCCGCGGCAGGTAAGTGCCAGGGAAT
GTTTGTTCTTAAATACCATCGCTCCAGGGAATGTTTGTTCTTAAATACCATCT
ACTGACACTGACATCCACTTTTTCTTTTTCTCCACAGGTATCGATCCACCATGC
AAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGCGATTCTGCTTTAGTG
CCACCAGAAGATACTACCTGGGTGCAGTGGAAGTGTCTATGGGACTATATGCA
AAGTGATCTCGGTGAGCTGCCTGTGGACGCAAGATTTCTCCTAGAGTGCCA
AAATCTTTTCCATTCAACACCTCAGTCGTGTACAAAAAGACTCTGTTTGTAGA
ATTACGGATCACCTTTTCAACATCGCTAAGCCAAGGCCACCCTGGATGGGTC
TGCTAGGTCCTACCATCCAGGCTGAGGTTTATGATACAGTGGTCATTACACTT
AAGAACATGGCTTCCCATCCTGTCAGTCTTCATGCTGTTGGTGTATCCTACTG
GAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAA
AGAAGATGATAAAGTCTTCCCTGGTGGAAGCCATACATATGTCTGGCAGGTC
CTGAAAGAGAATGGTCCAATGGCCTCTGACCCACTGTGCCTTACCTACTCATA
TCTTTCTCATGTGGACCTGGTAAAAGACTTGAATTCAGGCCTCATTGGAGCCC
TACTAGTATGTAGAGAAGGGAGTCTGGCCAAGGAAAAGACACAGACCTTGCA
CAAATTTATACTACTTTTTTGCTGTATTTGATGAAGGGAAAAGTTGGCACTCAG
AAACAAAGAACTCCTTGATGCAGGATAGGGATGCTGCATCTGCTCGGGCCTG
GCCTAAAATGCACACAGTCAATGGTTATGTAAACAGGTCTCTGCCAGGTCTG
ATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTGGAATGGGCACCA
CTCCTGAAGTGCACCTCAATATTCCTCGAAGGTCACACATTTCTTGTGAGGAAC
CATCGCCAGGCGTCCTTGGAATCTCGCCAATAACTTTCTTACTGCTCAAAC
ACTCTTGATGGACCTTGGACAGTTTCTACTGTTTTGTATATCTCTTCCCACCA
ACATGATGGCATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCC
CAACTACGAATGAAAAATAATGAAGAAGCGGAAGACTATGATGATGATCTTA
CTGATTCTGAAATGGATGTGGTCAGGTTTGATGATGACAACTCTCCTTCTTT
ATCCAAATTCGCTCAGTTGCCAAGAAGCATCCTAAAACCTGGGTACATTACAT
TGCTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTCCTCGCCCCCGATG
ACAGAAGTTATAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAG
GAAGTACAAAAAAGTCCGATTTATGGCATAACACAGATGAAACCTTTAAGACT
CGTGAAGCTATTCAGCATGAATCAGGAATCTTGGGACCTTTACTTTATGGGG
AAGTTGGAGACACACTGTTGATTATATTTAAGAATCAAGCAAGCAGACCATA
TAACATCTACCCTCACGGAATCACTGATGTCCGTCCTTTGTATTCAAGGAGAT
TACCAAAGGTGTAAAACATTTGAAGGATTTTCCAATTCTGCCAGGAGAAAT
ATTCAAATATAAATGGACAGTGAAGTGTAGAAGATGGGCCAACTAAATCAGAT
CCTCGGTGCCTGACCCGCTATTACTCTAGTTTCGTTAATATGGAGAGAGATCT
AGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATCTGTAGATC
AAAGAGGAAACCAGATAATGTCAGACAAGAGGAATGTCATCCTGTTTTCTGT
ATTTGATGAGAACCGAAGCTGGTACCTCACAGAGAATATACAACGCTTTCTC

CCCAA.TCCAGCTGGAGTGCAGCTTGAGGATCCAGAGTTCCAAGCCTCCAACA
TCATGCACAGCATCAATGGCTATGTTTTTGATAGTTTGCAGTTGTCAGTTTGT
TTGCATGAGGTGGCATACTGGTACATTCTAAGCATTGGAGCACAGACTGACT
TCCTTTCTGTCTTCTTCTCTGGATATACCTTCAAACACAAAATGGTCTATGAA
GACACACTCACCTATTCCCATTCTCAGGAGAAACTGTCCTCATGTGCGATGGA
AAACCCAGGTCTATGGATTCTGGGGTGCCACAACCTCAGACTTTCGGAACAGA
GGCATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACAACCTGGTGATT
ATTACGAGGACAGTTATGAAGATATTTACGCATACTTGCTGAGTAAAAACAA
TGCCATTGAACCAAGAAGCTTCTCCCAGAATCCACCAGTCTTGAAACGCCATC
AACGCGAAATAAECTCGTACTACTCTTCAGTCAGATCAAGAGGAAATTGACTA
TGATGATAACCATATCAGTTGAAATGAAGAAGGAAGATTTTGACATTTATGAT
GAGGATGAAAATCAGAGCCCCCGCAGCTTTCAAAGAAAACACGACACTATT
TTATTGCTGCAGTGGAGAGGCTCTGGGATTATGGGATGAGTAGCTCCCCACA
TGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAAAGTT
GTTTTCCAGGAATTTACTGATGGCTCCTTTACTCAGCCCTTATACCGTGGAGA
ACTAAATGAACATTTGGGACTCCTGGGGCCATATATAAGAGCAGAAGTTGAA
GATAATATCATGGTAACCTTCAGAAATCAGGCCTCTCGTCCCTATTCTTCTA
TTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGAGCAGAACCTAGA
AAAAACTTTGTCAAGCCTAATGAAACCAAACTTACTTTTGAAAGTGCAAC
ATCATATGGCACCCACTAAAGATGAGTTTGACTGCAAAGCCTGGGCTTATTTCT
TCTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCT
GGTCTGCCACACTAACACACTGAACCTGCTCATGGGAGACAAGTGACAGTA
CAGGAATTTGCTCTGTTTTTACCATCTTTGATGAGACCAAAAGCTGGTACTT
CACTGAAAATATGGAAAGAACTGCAGGGCTCCCTGCAATATCCAGATGGAA
GATCCCACTTTTAAAGAGAATTATCGCTTCCATGCAATCAATGGCTACATAAT
GGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTGATGGTAT
CTGCTCAGCATGGGCAGCAATGAAAACATCCATTCTATTCAATTCAGTGGACA
TGTGTTCACTGTACGAAAAAAGAGGAGTATAAAATGGCACTGTACAATCTC
TATCCAGGTGTTTTTGAGACAGTGGAAATGTTACCATCCAAAGCTGGAATTTG
GCGGGTGGAATGCCTTATTGGCGAGCATCTACATGCTGGGATGAGCACACTT
TTTCTGGTGTACAGCAATAAGTGTGCACTCCCCTGGGAATGGCTTCTGGACA
CATTAGAGATTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCA
AAGCTGGCCAGACTTCATTATTCCGGATCAATCAATGCCTGGAGACCAAGG
AGCCCTTTTCTTGGATCAAGGTGGATCTGTTGGCACCAATGATTATTCACGGC
ATCAAGACCCAGGGTGCCCGTCAGAAGTTCTCCAGCCTCTACATCTCTCAGTT
TATCATCATGTATAGTCTTGATGGGAAGAAGTGGCAGACTTATCGAGGAAAT
TCCACTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTATCTGGGATAAA
ACACAATATTTTAAACCCTCCAATTATTGCTCGATACATCCGTTTGCACCCAA
CTCATTATAGCATTTCGCAGCACTCTTCGCATGGAGTTGATGGGCTGTGATTTA
AATAGTTGCAGCATGCCATTGGGAATGGAGAGTAAAGCAATATCAGATGCAC
AGATTACTGCTTCATCCTACTTTACCAATATGTTTGCCACCTGGTCTCCTTCAA
AAGCTCGACTTCACCTCCAAGGGAGGAGTAATGCCTGGAGACCTCAGGTGAA
TAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGAAAGTCACA
GGAGTAACTACTCAGGGAGTAAAATCTCTGCTTACCAGCATGTATGTGAAGG
AGTTCCCTCATCTCCAGCAGTCAAGATGGCCATCAGTGGACTCTCTTTTTTTCAG
AATGGCAAAGTAAAGGTTTTTCAGGGAAATCAAGACTCCTTCACACCTGTGG

TGAACTCTCTAGACCCACCGTTACTGACTCGCTACCTTCGAATTCACCCCCAG
AGTTGGGTGCACCAGATTGCCCTGAGGATGGAGGTTCTGGGCTGCGAGGCAC
AGGACCTCTACTGACTCGAGCCTAATAAAGGAAATTTATTTTCATTGCAATAG
TGTGTTGGTTTTTTGTGTGCGGCCGCAGGAACCCCTAGTGATGGAGTTGGCCA
CTCCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGGCGACCAAAGGTCGCC
CGACGCCCCGGGCTTTGCCCGGGCGGCCTCAGTGAGCGAGCGAGCGCGCAGCT
GCCTGCAGGACAT

FIGURE 7




FIGURE 8






FIGURE 9




BEST AVAILABLE COPY

FIGURE 10



BEST AVAILABLE COPY

FIGURE 11



BEST AVAILABLE COPY

SEQUENCE LISTING

<110> Couto, Linda B.
Colosi, Peter C.

5

<120> ADENO-ASSOCIATED VECTORS FOR EXPRESSION OF FACTOR VIII
BY TARGET
CELLS

10 <130> AVIGEN-03743

<140> not yet assigned
<141> 1999-07-30

15 <150> 60/125,974
<151> 1999-03-24

<150> 60/104,994
<151> 1998-10-20

20

<160> 14

<170> PatentIn Ver. 2.0

25 <210> 1
<211> 59
<212> DNA
<213> Artificial Sequence30 <220>
<223> Description of Artificial Sequence: Synthetic

<400> 1
cccaagcttg cggccgccc ggtgccgcc ctaggcaggt aagtgccgtg tgtgggtcc 59

35

<210> 2

<211> 59

<212> DNA

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 2

10 ccgctcgagc agagctctat ttgcatggtg gaatcgatgc cgcgggaacc acacacggc 59

<210> 3

<211> 103

<212> DNA

15 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

20 <400> 3

cccaagcttg cggccgccc ggtgccgcc ctaggcaggt aagtgccgtg tgtgggtccc 60
gcggcatcga ttccaccatg caaatagagc tctgctcgag cgg 103

<210> 4

25 <211> 57

<212> DNA

<213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence: Synthetic

<400> 4

ttcccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactga 57

35 <210> 5

<211> 57

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 5

gaatcgatac ctgtggagaa aaagaaaaag tggatgtcag tgtcagtaat tcaaggc 57

10

<210> 6

<211> 99

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 6

20 ttcccgcggg cctggcctct ttacgggtta tggcccttgc gtgccttgaa ttactgacac 60
tgacatccac tttttctttt tctccacagg tatcgattc 99

<210> 7

<211> 100

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

30

<400> 7

agggaatggt tggtcttaaa taccatccag ggaatgtttg ttcttaaata ccatccaggg 60
aatgtttggt cttaaatacc atctacagtt attggttaaa 100

35 <210> 8

<211> 59

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 8

ggaaaggtga tctgtgtgca gaaagactcg ctctaataata cttctttaac caataactg 59

10

<210> 9

<211> 144

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 9

20 agggaaatggt tgttcttaaa taccatccag ggaatgtttg ttcttaaata ccatccaggg 60
aatgtttgtt cttaaatacc atctacagtt attggttaaa gaagtatatt agagcgagtc 120
tttctgcaca cagatcacct ttcc 144

<210> 10

25 <211> 59

<212> DNA

<213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence: Synthetic

<400> 10

tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgc 59

35 <210> 11

<211> 59

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence: Synthetic

<400> 11

tcgagcggcc gcacacaaaa aaccaacaca cagatctcta gagctctgat cttttatct 59

10

<210> 12

<211> 63

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 12

20 tcgagaataa aagatcagag ctctagagat ctgtgtgttg gttttttgtg tgcggccgct 60
cga 63

<210> 13

<211> 11933

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic

30

<400> 13

cagctgcgcg ctgcctcgct cactgaggcc gcccgggcaa agcccgggcg tcgggcgacc 60
tttggctgcc cggcctcagt gagcgagcga gcgcgcagag agggagtggc caactccatc 120
actagggggtt cctgcggccg ccagagggaat gtttgttctt aaataccatc cagggaatgt 180
35 ttgttcttaa ataccatcca gggaatgttt gttcttaaata accatctaca gttattgggt 240

aaagaagtat attagagcga gtctttctgc acacagatca cctttccggg tgccgccctt 300
 aggcaggtaa gtgccgtgtg tggttcccg cggcctggcc tctttacggg ttatggccct 360
 tgcgtgcctt gaattactga cactgacatc cactttttct ttttctccac aggtatcgat 420
 tccaccatgc aaatagagct ctccacctgc ttctttctgt gccttttgcg attctgcttt 480
 5 agtgcacca gaagatacta cctgggtgca gtggaactgt catgggacta tatgcaaagt 540
 gatctcgggtg agctgcctgt ggacgcaaga tttcctccta gagtgcctaa atcttttcca 600
 ttcaacacct cagtcgtgta caaaaagact ctgtttgtag aattcacgga tcaccttttc 660
 aacatcgcta agccaaggcc accctggatg ggtctgctag gtcctacct ccaggctgag 720
 gtttatgata cagtgggtcat tacacttaag aacatggctt cccatcctgt cagtcttcat 780
 10 gctgttggtg tatectactg gaaagcttct gagggagctg aatatgatga tcagaccagt 840
 caaagggaga aagaagatga taaagtcttc cctgggtggaa gccatacata tgtctggcag 900
 gtectgaaag agaatgggtcc aatggcctct gaccactgt gccttaccta ctcatatctt 960
 tctcatgtgg acctggtaaa agacttgaat tcaggcctca ttggagccct actagtatgt 1020
 agagaagggg gtctggccaa ggaaaagaca cagaccttgc acaaatttat actacttttt 1080
 15 gctgtatttg atgaagggaa aagttggcac tcagaaacaa agaactcctt gatgcaggat 1140
 agggatgctg catctgctcg ggcctggcct aaaatgcaca cagtcaatgg ttatgtaaac 1200
 aggtctctgc caggtctgat tggatgccac aggaaatcag tctattggca tgtgattgga 1260
 atgggcacca ctctgaagt gcactcaata ttcctcgaag gtcacacatt tcttgtaggg 1320
 aaccatcgcc aggcgtcctt ggaaatctcg ccaataactt tccttactgc tcaaactc 1380
 20 ttgatggacc ttggacagtt tctactgttt tgcataatct cttcccacca acatgatggc 1440
 atggaagctt atgtcaaagt agacagctgt ccagaggaac cccaactacg aatgaaaaat 1500
 aatgaagaag cggaagacta tgatgatgat ctactgatt ctgaaatgga tgtggtcagg 1560
 tttgatgatg acaactctcc ttcctttatc caaattcgct cagttgccaa gaagcactct 1620
 aaaacttggg tacattacat tgctgctgaa gaggaggact gggactatgc tcccttagtc 1680
 25 ctgcgccccg atgacagaag ttataaaagt caatatttga acaatggccc tcagcggatt 1740
 ggtaggaagt acaaaaaagt ccgatttatg gcatacacag atgaaacctt taagactcgt 1800
 gaagctattc agcatgaatc aggaatcttg ggacctttac tttatgggga agttggagac 1860
 aactgttgta ttatatttaa gaatcaagca agcagacct ataacatcta ccctcacgga 1920
 atcactgatg tccgtccttt gtattcaagg agattaccaa aaggtgtaaa acatttgaag 1980
 30 gattttccaa ttctgccagg agaaatattc aaatataaat ggacagtgc tgtagaagat 2040
 gggccaacta aatcagatcc tcggtgctg acccgctatt actctagttt cgtaaatatg 2100
 gagagagatc tagcttcagg actcattggc cctctcctca tctgctacaa agaactctgta 2160
 gatcaaagag gaaaccagat aatgtcagac aagaggaatg tcacctgtt ttctgtattt 2220
 gatgagaacc gaagctggta cctcacagag aatatacaac gctttctccc caatccagct 2280
 35 ggagtgcagc ttgaggatcc agagttccaa gcctccaaca tcatgcacag catcaatggc 2340

tatgtttttg atagtttgca gttgtcagtt tgtttgcatg aggtggcata ctggtacatt 2400
 ctaagcattg gagcacagac tgacttcctt tctgtcttct tctctggata taccttcaaa 2460
 cacaaaatgg tctatgaaga cacactcacc ctattcccat tctcaggaga aactgtcttc 2520
 atgtcgatgg aaaacccagg tctatggatt ctgggggtgcc acaactcaga ctttcggaac 2580
 5 agagggcatga ccgccttact gaagggttct agttgtgaca agaactctgg tgattattac 2640
 gaggacagtt atgaagatat ttcagcatat ttgctgagta aaaacaatgc cattgaacca 2700
 agaagcttcg aaataactcg tactactctt cagtcagatc aagaggaaaat tgactatgat 2760
 gataccatat cagttgaaat gaagaaggaa gattttgaca tttatgatga ggatgaaaat 2820
 cagagccccc gcagctttca aaagaaaaca cgacactatt ttattgctgc agtggagagg 2880
 10 ctctgggatt atgggatgag tagctcccca catgttctaa gaaacagggc tcagagtggc 2940
 agtgtccctc agttcaagaa agttgttttc caggaattta ctgatggctc ctttactcag 3000
 cccttatacc gtggagaact aaatgaacat ttgggactcc tggggccata tataagagca 3060
 gaagttgaag ataatatcat ggtaactttc agaaatcagg cctctcgtcc ctattccttc 3120
 tattctagcc ttatttctta tgaggaagat cagaggcaag gagcagaacc tagaaaaaac 3180
 15 tttgtcaagc ctaatgaaac caaaacttac ttttggaag tgcaacatca tatggcacc 3240
 actaaagatg agtttgactg caaagcctgg gcttatttct ctgatgttga cctggaaaaa 3300
 gatgtgcact caggcctgat tggaccctt ctggtctgcc aactaacac actgaaccct 3360
 gctcatggga gacaagtgc agtacaggaa tttgctctgt ttttcacat ctttgatgag 3420
 accaaaagct ggtacttcac tgaaaatat gaaagaaact gcagggtcc ctgcaatatt 3480
 20 cagatggaag atcccacttt taaagagaat tatcgcttcc atgcaatcaa tggctacata 3540
 atggatacac tacctggctt agtaatggct caggatcaaa ggattcgatg gtatctgctc 3600
 agcatgggca gcaatgaaaa catccattct attcatttca gtggacatgt gttcactgta 3660
 cgaaaaaaag aggagtataa aatggcactg tacaatctct atccagggtg ttttgagaca 3720
 gtggaaatgt taccatccaa agctggaatt tggcggtgg aatgccttat tggcgagcat 3780
 25 ctacatgctg ggatgagcac actttttctg gtgtacagca ataagtgtca gactcccctg 3840
 ggaatggctt ctggacacat tagagatttt cagattacag cttcaggaca atatggacag 3900
 tgggcccaca agctggccag acttcattat tccggatcaa tcaatgcctg gagcaccaag 3960
 gagccctttt cttggatcaa ggtggatctg ttggcaccaa tgattattca cggcatcaag 4020
 acccaggggtg ccgctcagaa gttctccagc ctctacatct ctcagtttat catcatgtat 4080
 30 agtcttgatg ggaagaagtg gcagacttat cgaggaaatt ccactggaac cttaatggctc 4140
 ttctttggca atgtggattc atctgggata aaacacaata tttttaacct tccaattatt 4200
 gctcgataca tccgtttgca cccaactcat tatagcattc gcagcactct tcgcatggag 4260
 ttgatgggct gtgatttaaa tagttgcagc atgccattgg gaatggagag taaagcaata 4320
 tcagatgcac agattactgc ttcattctac tttaccaata tgtttgccac ctggtctcct 4380
 35 tcaaaagctc gacttcacct ccaagggagg agtaatgcct ggagacctca ggtgaataat 4440

ccaaaagagt ggctgcaagt ggacttccag aagacaatga aagtcacagg agtaactact 4500
 cagggagtaa aatctctgct taccagcatg tatgtgaagg agttcctcat ctccagcagt 4560
 caagatggcc atcagtggac tctctttttt cagaatggca aagtaaagggt ttttcagggga 4620
 aatcaagact ccttcacacc tgtggtgaac tctctagacc caccgttact gactcgctac 4680
 5 cttcgaattc acccccagag ttgggtgcac cagattgccc tgaggatgga gggtctgggc 4740
 tgcgaggcac aggacctcta ctgactcgag aataaaagat cagagctcta gagatctgtg 4800
 tgttgggtttt ttgtgtgcgg ccgcaggaac ccctagtgat ggagttggcc actccctctc 4860
 tgcgcgctcg ctcgctcact gaggccgggc gaccaaagggt cgcccgacgc ccgggctttg 4920
 cccggggcggc ctcagtgagc gagcgagcgc gcagctgcct gcaggacatg tgagcaaaag 4980
 10 gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcggtttttc cataggctcc 5040
 gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aaccgcagac 5100
 gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct cctgttccga 5160
 ccctgccgct taccggatac ctgtccgctt ttctcccttc gggaaagcgtg gcgctttctc 5220
 atagctcacg ctgtaggtat ctgagttcgg tgtaggtcgt tcgctccaag ctgggctgtg 5280
 15 tgcacgaacc ccccggtcag ccgaccgct gcgccttacc cggtaactat cgtcttgagt 5340
 ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca 5400
 gagcgaggta tgtaggcgggt gctacagagt tcttgaagtg gtggcctaac tacggctaca 5460
 ctagaaggac agtatttggt atctgcgctc tgctgaagcc agttaccttc ggaaaaagag 5520
 ttggtagctc ttgatccggc aaacaaacca ccgctggtag cggtaggtttt tttgtttgca 5580
 20 agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc ttttctacgg 5640
 ggtctgacgc tcagtggaaac gaaaactcac gttaagggat tttggtcatg agattatcaa 5700
 aaaggatctt cacctagatc cttttaaatt aaaaatgaag ttttaaataca atctaaagta 5760
 tatatgagta aacttggtct gacagttacc aatgcttaat cagtgaggca cctatctcag 5820
 cgatctgtct atttcgttca tccatagttg cctgactccc cgtcgtgtag ataactacga 5880
 25 tacgggaggg cttaccatct ggccccagtg ctgcaatgat accgcgagac ccacgctcac 5940
 cggctccaga tttatcagca ataaaccagc cagccggaag ggccgagcgc agaagtggtc 6000
 ctgcaacttt atccgcctcc atccagtcta ttaattggtg ccgggaagct agagtaagta 6060
 gttcgccagt taatagtttg cgcaacggtg ttgccattgc tacaggcatc gtggtgtcac 6120
 gctcgctggt tggatgggt tcatcagct ccggttccca acgatcaagg cgagttacat 6180
 30 gatecccat gttgtgcaaa aaagcgggta gtccttcgg tcctccgatc gttgtcagaa 6240
 gtaagttggc cgcagtgtta tcaactcatg ttatggcagc actgcataat tctcttactg 6300
 tcatgccatc cgtaagatgc ttttctgtga ctggtgagta ctcaaccaag tcatctgag 6360
 aatagtgtat gggcgaccg agttgctctt gcccggcgtc aatacgggat aataccgcgc 6420
 cacatagcag aactttaaaa gtgctcatca ttggaaaacg ttcttcgggg cgaaaactct 6480
 35 caaggatctt accgctgttg agatccagtt cgatgtaacc cactcgtgca cccaactgat 6540

cttcagcatc ttttactttc accagcgttt ctgggtgagc aaaaacagga aggcaaaatg 6600
 ccgcaaaaaa gggaataagg gcgacacgga aatgttgaat actcatactc ttcttttttc 6660
 aatattattg aagcatttat caggggttatt gtctcatgag cggatacata tttgaatgta 6720
 tttagaaaaa taaacaaata ggggttccgc gcacatttcc ccgaaaagtg ccacctgacg 6780
 5 tctaagaaac cattattatc atgacattaa cctataaaaa taggcgtatc acgaggccct 6840
 ttcgctctcg gcgtttcggg gatgacgggtg aaaacctctg acacatgcag ctcccggaga 6900
 cggtcacagc ttgtctgtaa gcggatgccg ggagcagaca agcccgtcag ggcgcgtcag 6960
 cgggtgttgg cgggtgtcgg ggctggctta actatgcggc atcagagcag attgtactga 7020
 gagtgcacca taaaattgta aacgttaata ttttggttaa attcgcgtta aatttttgtt 7080
 10 aaatcagctc attttttaac caataggccg aaatcggcaa aatcccttat aaatcaaaag 7140
 aatagcccga gataggggtg agtgttgttc cagtttgga caagagtcca ctattaaaga 7200
 acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg 7260
 aaccatcacc caaatcaagt tttttggggc cgaggtgccg taaagcacta aatcggaacc 7320
 ctaaagggag cccccgattt agagcttgac ggggaaagcc ggcgaacgtg gcgagaaagg 7380
 15 aagggagaa agcgaaagga gcgggcgcta gggcgctggc aagtgtagcg gtcacgctgc 7440
 gcgtaaccac cacaccgcc gcgcttaatg cgccgctaca gggcgctac tatggttgc 7500
 ttgacgtatg cgggtgtgaa taccgcacag atgcgtaagg agaaaatacc gcatcaggcc 7560
 gtaacctgtc ggatcaccgg aaaggacccg taaagtata atgattatca tctacatc 7620
 acaacgtgcg tggaggccat caaacacgt caaataatca attatgacgc aggtatcgta 7680
 20 ttaattgatc tgcataact taacgtaaaa acaacttcag acaatacaaa tcagcgacac 7740
 tgaatacggg gcaacctcat gtcaacgaag aacagaacct gcagaacaac aaccgcaac 7800
 atccgctttc ctaaccaa atgattgaaca attaatatcg ctcttgagca aaaaggggtc 7860
 ggggaatttct cagcctgggt cattgaagcc tgccgtcgga gactaacgtc agaaaagaga 7920
 gcatatacat caattaaaag tgatgaagaa tgaacatccc gcgttcttcc ctccgaacag 7980
 25 gacgatattg taaattcact taattacgag ggcattgcag taattgagtt gcagttttac 8040
 cactttctct acagtgcag actgcgtgtt ggctctgtca cagactaaat agtttgaatg 8100
 attagcagtt atggtgatca gtcaaccacc agggataat ccttcatatt attatcgtgc 8160
 ttcaccaacg ctgcctcaat tgctctgaat gcttccagag acaccttatg ttctatacat 8220
 gcaattacaa catcagggtg actcatagaa atggtgctat taagcatatt ttttacacga 8280
 30 atcagatcca cggagggatc atcagcagat tgttctttat tcattttgtc gctccatgcg 8340
 cttgctcttc atctagcggg taaaatatta cttcaaatct ttctgtatga agatttgagc 8400
 acgttggcct tacatacatc tgtcgggtgt atttccctcc agaatgccag caggaccgca 8460
 ctttgttacg caaccaatac tattaagtga aaacattcct aatatttgac ataaatcatc 8520
 aacaaaacac aaggagggtc gaccagattg aaacgataaa aacgataatg caaactacgc 8580
 35 gccctcgtat cacatggaag gttttaccaa tggctcaggt tgccattttt aaagaaatat 8640

tcgatcaagt gcgaaaagat ttagactgtg aattgtttta ttctgaacta aaacgtcaca 8700
 acgtctcaca ttatatattac tatctagcca cagataatat tcacatcgtg ttagaaaacg 8760
 ataacaccgt gttaataaaa ggacttaaaa aggttgtaaa tgttaaattc tcaagaaaca 8820
 cgcattcttat agaaacgtcc tatgataggt tgaaatcaag agaaatcaca ttctagcaat 8880
 5 acaggggaaaa tcttgctaaa gcaggagttt tccgatgggt taaaaatc catgaacata 8940
 aaagatatata ctataccttt gataattcat tactatttac tgagagcatt cagaacacta 9000
 caaaaatctt tccacgctaa atcataacgt ccggttttctt ccgtgtcagc accggggcgt 9060
 tggcataatg caatacgtgt acgcgctaaa ccctgtgtgc atcgttttaa ttattcccgg 9120
 aactcccgc agagaagttc cccgtcaggg ctgtggacat agttaatccg ggaatacaat 9180
 10 gacgattcat cgcacctgac atacattaat aaatattaac aatatgaaat ttcaactcat 9240
 tgttttaggg ttgtttaatt ttctacacat acgattctgc gaacttcaaa aagcatcggg 9300
 aataacacca tgaaaaaaat gctactcgct actgcgctgg ccctgcttat tacaggatgt 9360
 gctcaacaga cgtttactgt tcaaaacaaa ccggcagcag tagcaccaaa ggaaaccatc 9420
 acccatcatt tcttcgtttc tggaattggg cagaagaaaa ctgtcgatgc agccaaaatt 9480
 15 tgtggcgggc cagaaaaatgt tgtaaaaaca gaaaccagc aaacattcgt aaatggattg 9540
 ctcggtttta ttactttagg catttatact ccgctggaag cgcgtgtgta ttgctcacia 9600
 taattgcatg agttgcccac cgcgatatgg gcaactctat ctgcactgct cattaatata 9660
 cttctgggtt ccttccagtt gtttttgcac agtgatcagc ctctctctga gggtgaaata 9720
 atcccgttca gcggtgtctg ccagtcgggg ggaggctgca ttatccacgc cggaggcggg 9780
 20 ggtggcttca cgcactgact gacagactgc tttgatgtgc aaccgacgac gaccagcggc 9840
 aacatcatca cgcagagcat cattttcagc tttagcatca gctaactcct tcgtgtattt 9900
 tgcacgagc gcagcaacat cacgctgacg catctgcatg tcagtaattg ccgcgttcgc 9960
 cagcttcagt tctctggcat tttgtcgcg ctgggctttg taggtaattg cgttatcacg 10020
 gtaatgatta acagcccatg acaggcagac gatgatgcag ataaccagag cggagataat 10080
 25 cgcggtgact ctgctcatac atcaatctct ctgaccgttc cggccgcttc tttgaatttt 10140
 gcaatcaggc tgtcagcctt atgctcgaac tgaccataac cagcgcccgg cagtgaagcc 10200
 cagatattgc tgcaacggtc gattgcctga cggatatcac cacgatcaat cataggtaaa 10260
 gcgccacgct ccttaatctg ctgcaatgcc acagcgtcct gacttttcgg agagaagtct 10320
 ttcaggccaa gctgcttgcg gtaggcatcc caccaacggg aaagaagctg gtagcgctccg 10380
 30 gcgcctgttg atttgagttt tgggttttagc gtgacaagtt tgcgagggtg atcggagtaa 10440
 tcagtaaata gctctccgcc tacaatgacg tcataacat gatttctggt tttctgacgt 10500
 ccgttatcag ttccctccga ccacgccagc atatcgagga acgccttacg ttgattattg 10560
 atttctacca tcttctactc cggctttttt agcagcgaag cgtttgataa gcgaaccaat 10620
 cgagtcagta ccgatgtagc cgataaacac gctcgttata taagcgagat tgctacttag 10680
 35 tccggcgaag tcgagaaggt cacgaatgaa ccaggcgata atggcgcaac tcgttgcgctc 10740

gattactgtt tttgtaaacg caccgccatt atatctgccg cgaaggtacg ccattgcaaa 10800
 cgcaaggatt gccccgatgc cttgttcctt tgccgcgaga atggcgggca acagggtcatg 10860
 tttttctggc atcttcatgt cttaccccca ataaggggat ttgctctatt taattaggaa 10920
 taaggtcgat tactgataga acaaatccag gctactgtgt ttagtaatca gatttgttcg 10980
 5 tgaccgatat gcacgggcaa aacggcagga gggtgttagc gcgacctcct gccacccgct 11040
 ttcacgaagg tcatgtgtaa aaggccgcag cgtaactatt actaatgaat tcaggacaga 11100
 cagtggctac ggctcagttt ggggtgtgct gttgctgggc ggcgatgacg cctgtacgca 11160
 tttggtgatc cggttctgct tccggtattc gcttaattca gcacaacgga aagagcactg 11220
 gctaaccagg ctccgcgact cttcacgatt atcgactcaa tgctcttacc tggtgtgcag 11280
 10 atataaaaaa tcccgaacc gttatgcagg ctctaactat tacctgcgaa ctgtttcggg 11340
 attgcatttt gcagacctct ctgcctgcga tgggtggagt tccagacgat acgtcgaagt 11400
 gaccaactag gcggaatcgg tagtaagcgc cgctctttt catctcacta ccacaacgag 11460
 cgaattaacc catcgttgag tcaaatttac ccaattttat tcaataagtc aatatcatgc 11520
 cgттаататг ttgccatccg tggcaatcat gctgctaacg tgtgaccgca ttcaaaatgt 11580
 15 tgtctgcgat tgactcttct ttgtggcatt gcaccaccag agcgtcatac agcggttaa 11640
 cagtgcgtga ccagggtgggt tgggtaaggt ttgggattag catcgtcaca gcgcgatatg 11700
 ctgcgcttgc tggcatcctt gaatagccga cgcctttgca tcttcgcac tctttctcga 11760
 caactctccc ccacagctct gttttggcaa tatcaaccgc acggcctgta ccatggcaat 11820
 ctctgcatct tgccccggc gtcgcggcac tacggcaata atccgcataa gcgaatgttg 11880
 20 cgagcacttg cagtacctt gccttagtat ttccttcaag ctgccccctgc agg 11933

<210> 14

<211> 4999

<212> DNA

25 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

30 <400> 14

cgccccctgca ggcagctgcg cgctcgtcgc ctactgagg ccgcccgggc aaagcccggg 60
 cgtcgggcca cttttggtcg cccggcctca gtgagcgagc gagcgcgag agagggagtg 120
 gccaaactcca tctactagggg ttcctgcygc cgcacgcgtg gtggcgcggg gtaaaactggg 180
 aaagtgatgt cgtgtactgg ctccgccttt tccccgaggg tgggggagaa ccgtatataa 240
 35 gtgcagtagt gcctgtgaac gttctttttc gcaacggggt tgccgccccg cggcaggtaa 300

gtgccagggga atgtttgttc ttaaatacca tcgctccagg gaatgtttgt tcttaaatac 360
 catctactga cactgacatc cactttttct ttttctccac aggtatcgat ccaccatgca 420
 aatagagctc tccacctgct tctttctgtg ccttttgcca ttctgcttta gtgccaccag 480
 aagatactac ctgggtgcag tggaaactgtc atgggactat atgcaaagtg atctcgggtga 540
 5 gctgcctgtg gacgcaagat ttctctctag agtgccaaaa tcttttccat tcaacacctc 600
 agtcgtgtac aaaaagactc tgttttaga attcacggat caccttttca acatcgctaa 660
 gccaaaggcca ccctggatgg gtctgctagg tctaccatc caggctgagg tttatgatac 720
 agtggtcatt acacttaaga acatggcttc ccatcctgtc agtcttcctg ctgttggtgt 780
 atcctactgg aaagcttctg agggagctga atatgatgat cagaccagtc aaagggagaa 840
 10 agaagatgat aaagtcttcc ctgggtggaag ccatacatat gtctggcagg tctgaaaga 900
 gaatgggtcca atggcctctg acccactgtg ccttacctac tcatatcttt ctcatgtgga 960
 cctggtaaaa gacttgaatt caggcctcat tggagcccta ctagtatgta gagaagggag 1020
 tctggccaag gaaaagacac agaccttgca caaatttata ctactttttg ctgtatttga 1080
 tgaagggaaa agttggcact cagaaacaaa gaactccttg atgcaggata gggatgctgc 1140
 15 atctgctcgg gcctggccta aaatgcacac agtcaatggg tatgtaaaca ggtctctgcc 1200
 aggtctgatt ggatgccaca ggaaatcagt ctattggcat gtgattggaa tgggcaccac 1260
 tctgaagtgc cactcaatat tctcgaagg tcacacattt cttgtgagga accatcgcca 1320
 ggcgtccttg gaaatctcgc caataacttt ccttactgct caaacactct tgatggacct 1380
 tggacagttt ctactgtttt gtcatatctc ttcccacaa catgatggca tggaaactta 1440
 20 tgtcaaagta gacagctgtc cagaggaacc ccaactacga atgaaaaata atgaagaagc 1500
 ggaagactat gatgatgatc ttactgattc tgaaatggat gtgggtcagg ttgatgatga 1560
 caactctcct tcctttatcc aaattcgctc agttgccaag aagcatccta aaacttgggt 1620
 acattacatt gctgctgaag aggaggactg ggactatgct cccttagtcc tcgccccga 1680
 tgacagaagt tataaaagtc aatatttgaa caatggccct cagcggattg gtaggaagta 1740
 25 caaaaaagtc cgatttatgg catacacaga tgaaaccttt aagactcgtg aagctattca 1800
 gcatgaatca ggaatcttgg gacctttact ttatggggaa gttggagaca cactgttgat 1860
 tatatttaag aatcaagcaa gcagaccata taacatctac cctcacggaa tctatgatgt 1920
 ccgtcctttg tattcaagga gattacaaaa aggtgtaaaa catttgaagg attttccaat 1980
 tctgccagga gaaatattca aatataaatg gacagtgact gtagaagatg ggccaactaa 2040
 30 atcagatcct cgggtgcctga cccgtatta ctctagtctt gttaatatgg agagagatct 2100
 agcttcagga ctcatgggcc ctctcctcat ctgctacaaa gaatctgtag atcaaagagg 2160
 aaaccagata atgtcagaca agaggaatgt catcctgttt tctgtatttg atgagaaccg 2220
 aagctggtac ctcacagaga atatacaacg ctttctcccc aatccagctg gagtgcagct 2280
 tgaggatcca gagttccaag cctccaacat catgcacagc atcaatggct atgtttttga 2340
 35 tagtttgcag ttgtcagttt gtttgcata ggtggcatac tggtagattc taagcattgg 2400

agcacagact gacttccttt ctgtcttctt ctctggatat accttcaaac acaaaatggt 2460
 ctatgaagac acactcacc ctttccatt ctccaggagaa actgtcttca tgtcgatgga 2520
 aaaccacaggt ctatggattc tgggggtgcc caactcagac tttcggaaca gaggcagatgac 2580
 cgccttactg aagggtttcta gttgtgacaa gaacactggt gattattacg aggacagtta 2640
 5 tgaagatatt tcagcactat tgctgagtaa aaacaatgcc attgaaccaa gaagcttctc 2700
 ccagaatcca ccagtcttga aacgccatca acgcgaaata actcgtacta ctcttcagtc 2760
 agatcaagag gaaattgact atgatgatac catatcagtt gaaatgaaga aggaagattt 2820
 tgacatttat gatgaggatg aaaatcagag ccccgagcagc tttcaaaaga aaacacgaca 2880
 ctattttatt gctgcagtgg agaggctctg ggattatggg atgagtagct ccccatagtc 2940
 10 tctaagaaac agggctcaga gtggcagtggt ccctcagttc aagaaagttg ttttcagga 3000
 atttactgat ggctccttta ctccagcctt ataccgtgga gaactaaatg aacatttggg 3060
 actcctgggg ccatatataa gagcagaagt tgaagataat atcatggtta ctttcagaaa 3120
 tcaggcctct cgtccctatt ccttctattc tagccttatt tcttatgagg aagatcagag 3180
 gcaaggagca gaacctagaa aaaactttgt caagcctaag gaaacaaaaa cttacttttg 3240
 15 gaaagtcaa catcatatgg caccactaa agatgagttt gactgcaaag cctgggctta 3300
 tttctctgat gttgacctgg aaaaagatgt gcactcaggg ctgattggac cccttctggt 3360
 ctgccacact aacacactga accctgctca tgggagacaa gtgacagtac aggaatttgc 3420
 tctgtttttc accatctttg atgagaccaa aagctggtac ttcactgaaa atatggaaa 3480
 aaactgcagg gctccctgca atatccagat ggaagatccc acttttaaaag agaattatcg 3540
 20 cttccatgca atcaatggct acataatgga taccactacct ggcttagtaa tggctcagga 3600
 tcaaaggatt cgatggtatc tgctcagcat gggcagcaat gaaaacatcc attctattca 3660
 tttcagtgga catgtgttca ctgtacgaaa aaaagaggag tataaaatgg cactgtacaa 3720
 tctctatcca ggtgtttttg agacagtgga aatgttacca tccaaagctg gaatttggcg 3780
 ggtggaatgc cttattggcg agcatctaca tgctgggatg agcacacttt ttctggtgta 3840
 25 cagcaataag tgtcagactc ccctgggaat ggcttctgga cacattagag attttcagat 3900
 tacagcttca ggacaatatg gacagtgggc cccaaagctg gccagacttc attattccgg 3960
 atcaatcaat gcctggagca ccaaggagcc cttttcttgg atcaagggtg atctgttggc 4020
 accaatgatt attcacggca tcaagacca ggggtgccgt cagaagttct ccagcctcta 4080
 catctctcag tttatcatca tgtatagtct tgatgggaag aagtggcaga cttatcgagg 4140
 30 aaattccact ggaaccttaa tggctcttct tggcaatgtg gattcatctg ggataaaaca 4200
 caatattttt aaccctccaa ttattgctcg atacatccgt ttgcacccaa ctcatatag 4260
 cattcgagc actcttcgca tggagttgat gggctgtgat ttaaatagtt gcagcatgcc 4320
 attgggaatg gagagtaaag caatatcaga tgcacagatt actgcttcat cctactttac 4380
 caatatgttt gccacctggt ctccctcaaa agctcgactt cacctccaag ggaggagtaa 4440
 35 tgccctggaga cctcaggtga ataatacaaa agagtggctg caagtggact tccagaagac 4500

aatgaaagtc acaggagtaa ctactcaggg agtaaaatct ctgcttacca gcatgtatgt 4560
gaaggagttc ctcactcca gcagtcaaga tggccatcag tggactctct tttttcagaa 4620
tggcaaagta aaggtttttc agggaaatca agactccttc acacctgtgg tgaactctct 4680
agaccaccg ttactgactc gctaccttcg aattcacccc cagagttggg tgcaccagat 4740
5 tgccctgagg atggaggttc tgggctgca ggcacaggac ctctactgac tcgagcctaa 4800
taaaggaaat ttattttcat tgcaatagtg tggttggttt ttgtgtgcgg ccgcaggaac 4860
ccctagtgat ggagttggcc actccctctc tgccgctcg ctcgctcact gaggccgggc 4920
gaccaaaggt cggccgacgc ccgggctttg cccgggcggc ctcagtgagc gagcgagcgc 4980
gcagctgcct gcaggacat 4999

10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24495

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 48/00; C12N 15/63, 15/00; C07H 21/04

US CL : 514/44; 435/320.1, 69.1; 424/93.6; 536/23.51

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44; 435/320.1, 69.1; 424/93.6; 536/23.51

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, CAPLUS, MEDLINE, BIOSIS, WEST

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ZATLOUKAL et al. In vivo production of human factor VIII in mice after intrasplenic implantation of primary fibroblasts transfected by receptor-mediated, adenovirus-augmented gene delivery. Proc. Natl. Acad. Sci., U.S.A. May 1994, Vol. 91, No. 11, pages 5148-5152, entire document.	1-17, 20-25, 26, 28
X	CONNELLY et al. Complete short-term correction of canine hemophilia A by in vivo gene therapy. Blood. 15 November 1996, Vol. 88, No. 10, pages 3846-3853, entire document.	1-30
X	CONNELLY et al. Sustained phenotypic correction of murine hemophilia A by in vivo gene therapy. Blood. 01 May 1998, Vol. 91, No. 9, pages 3273-3281, entire document.	1-30



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 DECEMBER 1999

Date of mailing of the international search report

10 FEB 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GAI (JENNIFER) MI LEE


Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24495

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SNYDER et al. Persistent and therapeutic concentrations of human factor IX in mice after hepatic gene transfer of recombinant AAV vector. Nat. Genet. July 1997, Vol. 16, No. 3, pages 270-276, entire document.	1-30
Y	LOZIER et al. Gene therapy and the hemophilias. JAMA, J. Am. Med. Assoc. 04 January 1994, Vol. 271, No. 1, pages 47-51, entire document.	1-30



THIS PAGE BLANK (USPTO)